

Datasheet for ABIN5008012

anti-OLIG1 antibody (AA 121-220) (AbBy Fluor® 750)[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	OLIG1
Binding Specificity:	AA 121-220
Reactivity:	Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This OLIG1 antibody is conjugated to AbBy Fluor® 750
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human OLIG1
Isotype:	IgG
Cross-Reactivity:	Rat
Predicted Reactivity:	Human,Mouse,Dog,Cow,Pig
Purification:	Purified by Protein A.

Target Details

Target:	OLIG1
Alternative Name:	OLIG1 (OLIG1 Products)

Target Details

Background:	Synonyms: Oligo 1, Oligo1, Oligodendrocyte lineage transcription factor 1, Oligodendrocyte specic bHLH transcription factor 1, Oligodendrocyte transcription factor 1. Basic domain helix loop helix protein class B 6, Basic domain helix loop helix protein class B6, BHLH B6, BHLHB 6, BHLHB6, bHLHe21, Class B basic helix-loop-helix protein 6, Class E basic helix-loop-helix protein 21, Olig 1, Olig1, OLIG1_HUMAN. Background: The oligodendrocyte lineage-specific basic helix-loop-helix (OLIG) family of transcription factors include OLIG1-OLIG3, which differ in tissue expression. OLIG1 and OLIG2 are specifically expressed in nervous tissue as gene regulators of oligodendrogenesis. OLIG2 is more widely expressed in embryonic brain than OLIG1, while OLIG3 is primarily expressed in non-neural tissues. OLIG1 and OLIG2 interact with the Nkx-2.2 homeodomain protein, which is responsible for directing ventral neuronal patterning in response to graded Sonic hedgehog signaling in the embryonic neural tube. These interactions between OLIG proteins and Nkx-2.2 appear to promote the formation of alternate cell types by inhibiting V3 interneuron development. OLIG1 and OLIG2 are abundantly expressed in oligodendroglioma and nearly absent in astrocytomas. Therefore, OLIG proteins are candidates for molecular markers of human glial brain tumors, which are the most common primary malignancies of the human brain.
-------------	--

Gene ID:	116448
----------	--------

UniProt:	Q8TAK6
----------	------------------------

Application Details

Application Notes:	IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200
--------------------	--

Restrictions:	For Research Use only
---------------	-----------------------

Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin

Handling

Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months