antibodies -online.com





anti-OPCML antibody (AA 101-200) (Alexa Fluor 680)



Go to Product page

()	1/0	r\ /1	014	
()	ve	I V I	-v	V

Quantity:	100 μL
Target:	OPCML
Binding Specificity:	AA 101-200
Reactivity:	Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This OPCML antibody is conjugated to Alexa Fluor 680
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human OPCML
Isotype:	IgG
Cross-Reactivity:	Rat
Predicted Reactivity:	Human,Mouse,Cow,Horse,Chicken,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	OPCML
Alternative Name:	OPCML (OPCML Products)

Target Details

Background:	Synonyms: opioid binding protein/cell adhesion molecule-like, OPCML, OBCAM, OPCM,	
	3732419F12, C230027C17, 2900075015RIK, Al844366, B930023M13RIK, Gm181, IGLON1,	
	MGC99974, OPCM_HUMAN.	
	Background: This gene encodes a member of the IgLON subfamily in the immunoglobulin	
	protein superfamily. The encoded protein is localized in the plasma membrane and may have	
	an accessory role in opioid receptor function. This gene has an ortholog in rat and bovine. The	
	opioid binding-cell adhesion molecule encoded by the rat gene binds opioid alkaloids in the	
	presence of acidic lipids, exhibits selectivity for mu ligands and acts as a GPI-anchored protein.	
	Since the encoded protein is highly conserved in species during evolution, it may have a	
	fundamental role in mammalian systems. Two transcript variants encoding different isoforms	
	have been found for this gene. [provided by RefSeq]	
Gene ID:	4978	
Application Details		
• •	IE(II IO D) 1.F0 200	
Application Notes:	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200 IF(ICC) 1:50-200	
	IF(IGG) 1.50°200	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and	
	50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be	
	handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	
Expiry Date:	12 months	