antibodies -online.com





anti-DPH1 antibody (AA 251-350) (Alexa Fluor 680)



Go to Product page

| \sim | | | |
|--------|-----|------|-----|
| | N/P | r\/I | i⊢₩ |

| Quantity: | 100 μL |
|----------------------|--|
| Target: | DPH1 |
| Binding Specificity: | AA 251-350 |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This DPH1 antibody is conjugated to Alexa Fluor 680 |
| Application: | Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |

Product Details

| Immunogen: | KLH conjugated synthetic peptide derived from human OVCA1 |
|-----------------------|---|
| Isotype: | IgG |
| Cross-Reactivity: | Human |
| Predicted Reactivity: | Mouse,Rat,Dog,Cow,Horse,Rabbit |
| Purification: | Purified by Protein A. |

Target Details

| Target: | DPH1 |
|-------------------|-----------------------|
| Alternative Name: | OVCA1 (DPH1 Products) |

Target Details

| Background: | Synonyms: Candidate tumor suppressor in ovarian cancer 1, Diphthamide biosynthesis protein | |
|---------------------|---|--|
| | 1, Diphthamide biosynthesis protein 2 homolog like 1, Diphthamide biosynthesis protein 2 like, | |
| | Diptheria toxin resistance protein required for diphthamide biosynthesis Saccharomyces like 1, | |
| | Diptheria toxin resistance protein required for diphthamide biosynthesis like 1, DPH like 1 S. | |
| | cerevisiae, DPH like 1, DPH1, DPH1 homolog S. cerevisiae, DPH1 homolog, DPH2 like 1, DPH2L, | |
| | DPH2L1, HsDph1, Ovarian cancer associated gene 1, Ovarian cancer associated gene 1 protein, | |
| | Ovarian tumor suppressor candidate 1, DPH1_HUMAN. | |
| | Background: OVCA1 is required for the first step in the synthesis of diphthamide, a unique | |
| | posttranslationally modified histidine found only in translation elongation factor-2 (EEF2, MIM | |
| | 130610). This modification is conserved from archaebacteria to humans and serves as the | |
| | target for ADP-ribosylation and inactivation of EEF2 by diphtheria toxin (DT) and Pseudomonas | |
| | exotoxin A. OVCA1 is one of several enzymes involved in synthesis of diphthamide in EEF2. | |
| Gene ID: | 1801 | |
| Application Details | | |
| Application Notes: | IF(IHC-P) 1:50-200 | |
| | IF(IHC-F) 1:50-200 | |
| | IF(ICC) 1:50-200 | |
| Restrictions: | For Research Use only | |
| Handling | | |
| Format: | Liquid | |
| Concentration: | 1 μg/μL | |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and | |
| | 50 % Glycerol. | |
| Preservative: | ProClin | |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be | |
| | handled by trained staff only. | |
| Storage: | -20 °C | |
| Storage Comment: | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles. | |
| Expiry Date: | 12 months | |
| | | |