antibodies -online.com





anti-ST3GAL6 antibody (C-Term)

2 Images



Go to Product page

Overview

0.1 mg
ST3GAL6
C-Term
Human, Mouse, Rat
Rabbit
Polyclonal
This ST3GAL6 antibody is un-conjugated
Western Blotting (WB), Enzyme Immunoassay (EIA), Immunofluorescence (IF)
ST3gal6 antibody was raised against a 16 amino acid peptide from near the carboxy terminus of Human ST3gal6.
IgG
This antibody detects ST3GAL6.
Species reactivity (tested):Human, Mouse and Rat
Species reactivity (tested):Human, Mouse and Rat Peptide Affinity Chromatography


Target Details

Background:	Sialyltransferases catalyze the transfer of sialic acid from cytidine 5-prime monophospho-N-
	acetylneuraminic acid (CMP-NeuAc) to terminal positions of glycoprotein and glycolipid
	carbohydrate groups. Terminal NeuAc residues are key determinants of carbohydrate
	structures, such as the sialyl-Lewis X determinants, and are widely distributed in many cell
	types. However, cancer cells often express more heavily sialylated glycans on their cell surface
	and this feature sometimes correlates with invasiveness. In contrast, expression of ST3gal6, a
	member of the sialyltransferase family that sialylates type II lactosamine structures on
	glycoproteins and glycolipids, was found to be significantly decreased by hypermethylation of
	the gene in gastrointestinal cancer. At least three isoforms of ST3gal6 are known to
	exist.Synonyms: 3-sialyltransferase, 3-sialyltransferase VI, CMP-NeuAc:beta-galactoside alpha-
	2, SIAT10, ST3Gal VI, ST3GalVI, Sialyltransferase 10, Type 2 lactosamine alpha-2
Gene ID:	10402
NCBI Accession:	NP_006091
Pathways:	Glycosaminoglycan Metabolic Process
Application Details	
Application Notes:	ELISA. Western blot: 1 - 2 μg/mL. Immunoflourescence.
	Other applications not tested.
	Optimal dilutions are dependent on conditions and should be determined by the user.
Restrictions:	For Research Use only
Handling	
Concentration:	1.0 mg/mL
Buffer:	PBS containing 0.02 % Sodium Azide as preservative
Preservative:	Sodium azide
Precaution of Use:	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which
	should be handled by trained staff only.
Handling Advice:	Avoid repeated freezing and thawing.
Storage:	4 °C/-20 °C
Storage Comment:	Store the antibody undiluted at 2-8 °C for one month or (in aliquots) at -20 °C for longer.



Immunofluorescence

Image 1. Immunocytochemistry of ST3gal6 in HeLa cells with this product at $5 \, \mu g/ml$.



Western Blotting

Image 2. Western blot analysis of ST3gal6 in HeLa cell lysate with this product at (A) 1 and (B) $2 \mu g/ml$.