# antibodies -online.com





# anti-P4HTM antibody (AA 201-300) (AbBy Fluor® 750)



Go to Product page

| ( ) | 11/0   | K\ /   | iew                 | 1 |
|-----|--------|--------|---------------------|---|
|     | $\cup$ | 'I V/I | $\square \vee \vee$ | ı |
|     |        |        |                     |   |

| Quantity:            | 100 μL   |  |
|----------------------|--|--|
| Target:              | P4HTM  |  |
| Binding Specificity: | AA 201-300   |  |
| Reactivity:          | Mouse, Rat   |  |
| Host:                | Rabbit   |  |
| Clonality:           | Polyclonal   |  |
| Conjugate:           | This P4HTM antibody is conjugated to AbBy Fluor® 750   |  |
| Application:         | Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |  |

## **Product Details**

| Immunogen:            | KLH conjugated synthetic peptide derived from human PH-4 |
|-----------------------|--|
| Isotype:              | IgG  |
| Cross-Reactivity:     | Mouse, Rat   |
| Predicted Reactivity: | Human,Dog,Cow,Pig,Horse,Rabbit                           |
| Purification:         | Purified by Protein A.                                   |

# **Target Details**

| Target:           | P4HTM                  |
|-------------------|------------------------|
| Alternative Name: | P4HTM (P4HTM Products) |

### **Target Details**

| Background: |
|-------------|
|-------------|

Synonyms: Transmembrane prolyl 4-hydroxylase, Proline hydrolase 4, proline 4-hydroxylase, P4htm, hypoxia-inducible factor prolyl 4-hydroxylase, P4H-TM, H-prolyl hydroxylase PH-4, P4HTM\_HUMAN, Hypoxia-inducible factor prolyl hydroxylase 4, H-PH4, H-prolyl hydroxylase 4, HPH-4, transmembrane prolyl 4-hydroxylase isoform a, P4HTM, EGLN4, HPH4, P4H-TM, PH-4, PHD4, P4H with transmembrane domain, Prolyl hydroxylase domain-containing 4, hypoxia inducible factor prolyl 4 hydroxylase, H-prolyl hydroxylase 4.

Background: The product of this gene belongs to the family of prolyl 4-hydroxylases. This protein is a prolyl hydroxylase that may be involved in the degradation of hypoxia-inducible transcription factors under normoxia. It plays a role in adaptation to hypoxia and may be related to cellular oxygen sensing. Alternatively spliced variants encoding different isoforms have been identified. [provided by RefSeq, Jul 2008].

# **Application Details**

Application Notes:

IF(IHC-P) 1:50-200

IF(IHC-F) 1:50-200

IF(ICC) 1:50-200

Restrictions:

For Research Use only

# Handling

| Format:            | Liquid   |
|--------------------|--|
| Concentration:     | 1 μg/μL  |
| Buffer:            | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.         |
| Preservative:      | ProClin  |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. |
| Storage:           | -20 °C   |
| Storage Comment:   | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.                                  |
| Expiry Date:       | 12 months  |