

Datasheet for ABIN500833
anti-STK39 antibody (C-Term)[Go to Product page](#)

2 Images

Overview

Quantity:	0.1 mg
Target:	STK39
Binding Specificity:	C-Term
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This STK39 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Enzyme Immunoassay (EIA)

Product Details

Immunogen:	Stk39 antibody was raised against a 19 amino acid peptide from near the carboxy terminus of human Stk39.
Isotype:	IgG
Specificity:	This antibody detects STK39 / SPAK at C-term.
Cross-Reactivity (Details):	Species reactivity (tested): Human, mouse
Purification:	Peptide affinity chromatography

Target Details

Target:	STK39
---------	-------

Target Details

Alternative Name:	STK39 / SPAK (STK39 Products)
Background:	<p>The serine/threonine kinase Stk39 belongs to the STE20 family, a group of kinases that are known to interact with inflammation-related kinases (such as p38, JNK, NKCC1, PKC-theta, WNK and MLCK), and with transcription factor AP-1. The STE 20 family is involved in diverse biological phenomena, including cell differentiation, cell transformation/ proliferation, cytoskeleton rearrangement, and the regulation of ion transporters. STK39 contains an N-terminal series of proline and alanine repeats (PAPA box), followed by a serine/threonine kinase catalytic domain and is abundantly expressed in the brain. STK39 is activated in response to hypotonic stress, leading to phosphorylation of several cation-chloride-coupled co-transporters. The catalytically active kinase specifically activates the p38 MAP kinase pathway, and its interaction with p38 decreases upon cellular stress, suggesting that this kinase may serve as an intermediate in the response to cellular stress. Recent studies show that STK39 tend to be a novel candidate gene for autism and hypertension. Synonyms: DCHT, STE20/SPS1-related proline-alanine-rich protein kinase, Serine/threonine-protein kinase 39, Ste-20-related kinase</p>
Gene ID:	27347
NCBI Accession:	NP_037365

Application Details

Application Notes:	<p>ELISA. Western blot: 1 - 2 µg/mL. Immunohistochemistry on paraffin sections.</p> <p>Other applications not tested.</p> <p>Optimal dilutions are dependent on conditions and should be determined by the user.</p>
Restrictions:	For Research Use only


Handling

Buffer:	PBS containing 0.02 % sodium azide
Preservative:	Sodium azide
Precaution of Use:	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Avoid repeated freezing and thawing.
Storage:	4 °C/-20 °C
Storage Comment:	Store at 2 - 8 °C for up to one month or (in aliquots) at -20 °C for longer.



Immunohistochemistry (Paraffin-embedded Sections)

Image 1. Immunohistochemistry of Stk39 in human brain tissue with this product at 2.5 µg/ml.



Western Blotting

Image 2. Western blot analysis of Stk39 in SK-N-SH cell lysate with this product at (A) 1 and (B) 2 µg/ml.