



[Go to Product page](#)

Datasheet for ABIN5008516

## anti-PCYT2 antibody (AA 315-389) (AbBy Fluor® 680)

### Overview

Quantity:	100 µL
Target:	PCYT2
Binding Specificity:	AA 315-389
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PCYT2 antibody is conjugated to AbBy Fluor® 680
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

### Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human PCYT2
Isotype:	IgG
Predicted Reactivity:	Human, Mouse, Rat, Dog, Cow, Sheep, Horse
Purification:	Purified by Protein A.

### Target Details

Target:	PCYT2
Alternative Name:	PCYT2 ( <a href="#">PCYT2 Products</a> )
Background:	Synonyms: CTP, phosphoethanolamine cytidyltransferase, ET, Ethanolamine-phosphate

## Target Details

---

cytidyltransferase, Ethanolaminephosphate Cytidyltransferase, PCY2, PCY2\_HUMAN, Pcyt2, Phosphate cytidyltransferase 2 ethanolamine, Phosphorylethanolamine transferase.

Background: Phosphatidylethanolamine (PtdEtn) is a major membrane phospholipid which serves to play a primary role in cell membrane structure and is also involved in cell division, cell signaling, activation, phagocytosis and autophagy. PCYT2 (Phosphorylethanolamine transferase), also known as Ethanolamine-phosphate cytidyltransferase, is a 389 amino acid protein that catalyzes the formation of CDP-ethanolamine from ethanolamine. This product combined with diacylglycerol form phosphatidylethanolamine via the de novo Kennedy pathway. PCYT2 is expressed at highest levels in heart, liver and skeletal muscle. Elevated levels of MyoD, reduced content of Sp1 and a changed ratio of Sp1 to Sp3 all together stimulate upregulation of PCTY2 transcription during C2C12 muscle cell differentiation. Disruption of the PCYT2 gene in mice leads to death after embryo implantation, establishing the necessity of PCYT2 for murine development.

## Application Details

---

Application Notes: IF(IHC-P) 1:50-200  
IF(IHC-F) 1:50-200  
IF(ICC) 1:50-200

Restrictions: For Research Use only

## Handling

---

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Expiry Date: 12 months