

Datasheet for ABIN5008559

anti-PDGFA antibody (AA 125-170) (AbBy Fluor® 680)



oo to rroudet page

| Overview | |
|-----------------------|---|
| Quantity: | 100 μL |
| Target: | PDGFA |
| Binding Specificity: | AA 125-170 |
| Reactivity: | Human, Mouse |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This PDGFA antibody is conjugated to AbBy Fluor® 680 |
| Application: | Western Blotting (WB), Flow Cytometry (FACS), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |
| Product Details | |
| Immunogen: | KLH conjugated synthetic peptide derived from human PDGF-A |
| Isotype: | IgG |
| Cross-Reactivity: | Human, Mouse |
| Predicted Reactivity: | Rat,Rabbit |
| Purification: | Purified by Protein A. |
| Target Details | |
| Target: | PDGFA |
| Alternative Name: | PDGF A (PDGFA Products) |

Target Details

| Target Details | | |
|---------------------|--|--|
| Background: | Synonyms: PDGF1, PDGF-A, Platelet-derived growth factor subunit A, PDGF subunit A, PDGF-1, | |
| | Platelet-derived growth factor A chain, Platelet-derived growth factor alpha polypeptide, PDGFA | |
| | Background: Growth factor that plays an essential role in the regulation of embryonic | |
| | development, cell proliferation, cell migration, survival and chemotaxis. Potent mitogen for cells | |
| | of mesenchymal origin. Required for normal lung alveolar septum formation during | |
| | embryogenesis, normal development of the gastrointestinal tract, normal development of | |
| | Leydig cells and spermatogenesis. Required for normal oligodendrocyte development and | |
| | normal myelination in the spinal cord and cerebellum. Plays an important role in wound healing. | |
| | Signaling is modulated by the formation of heterodimers with PDGFB (By similarity). | |
| Gene ID: | 5154 | |
| UniProt: | P04085 | |
| Pathways: | RTK Signaling, Fc-epsilon Receptor Signaling Pathway, EGFR Signaling Pathway, Neurotrophin | |
| | Signaling Pathway, Smooth Muscle Cell Migration, Platelet-derived growth Factor Receptor | |
| | Signaling | |
| Application Details | | |
| Application Notes: | FCM 1:20-100 | |
| | IF(IHC-P) 1:50-200 | |
| | IF(IHC-F) 1:50-200 | |
| | IF(ICC) 1:50-200 | |
| Restrictions: | For Research Use only | |
| Handling | | |
| Format: | Liquid | |
| Concentration: | 1 μg/μL | |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and | |
| | 50 % Glycerol. | |
| Preservative: | ProClin | |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be | |
| | handled by trained staff only. | |
| Storage: | -20 °C | |
| | | |

Handling

| Storage Comment: | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles. |
|------------------|---|
| Expiry Date: | 12 months |