



[Go to Product page](#)

Datasheet for ABIN5008721
anti-FNTA antibody (Alexa Fluor 680)

Overview

Quantity:	100 µL
Target:	FNTA
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This FNTA antibody is conjugated to Alexa Fluor 680
Application:	Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human PGGT1A/FNTA
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Purification:	Purified by Protein A.

Target Details

Target:	FNTA
Alternative Name:	PGGT1A (FNTA Products)
Background:	Synonyms: CAAX farnesyltransferase alpha subunit, Farnesyl protein transferase alpha subunit, Farnesyltransferase CAAX box alpha, Farnesyltransferase, CAAX box, alpha, FPTA, FTase alpha, GGTase I alpha, PGGT1A, Protein farnesyltransferase/geranylgeranyltransferase type I alpha subunit, Protein prenyltransferase alpha subunit repeat containing 2, PTAR2, Ras proteins

Target Details

prenyltransferase alpha, Ras proteins prenyltransferase subunit alpha, Type I protein geranyl geranyltransferase alpha subunit, FNTA_HUMAN.

Background: FNTA, also known as CAAX farnesyltransferase (FTase), attaches a farnesyl group from farnesyl pyrophosphate to cysteine residues at the fourth position from the C terminus of proteins that end in the so-called CAAX box, where C is cysteine, A is usually but not always an aliphatic amino acid, and X is typically methionine or serine. This type of posttranslational modification provides a mechanism for membrane localization of proteins that lack a transmembrane domain. This enzyme has the remarkable property of farnesylating peptides as short as four residues in length that conform to the CAAX consensus sequence. FNTA is also a specific cytoplasmic interactor of the transforming growth factor-beta and activin type I receptors. It is likely to be a key component of the signaling pathway which involves p21ras, an important substrate for farnesyltransferase.

Gene ID: 2339

Pathways: [Response to Water Deprivation, Regulation of G-Protein Coupled Receptor Protein Signaling](#)

Application Details

Application Notes: IF(IHC-P) 1:50-200

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Expiry Date: 12 months