antibodies - online.com







anti-CD248 antibody (C-Term)






\sim						
	1//	Д	r۱	/1	\triangle	٨

Quantity:	0.1 mg	
Target:	CD248	
Binding Specificity:	C-Term	
Reactivity:	Human, Mouse, Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This CD248 antibody is un-conjugated	
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunofluorescence (IF)	
Product Details		
lmmunogen:	TEM1 antibody was raised against a 14 amino acid peptide near the carboxy terminus of the human TEM1.	
Isotype:	IgG	
Specificity:	This antibody detects CD248 / TEM1 at C-term. It recognizes both isoforms.	
Cross-Reactivity (Details):	Species reactivity (tested):Human, Mouse, Rat	
Purification:	Peptide affinity chromatography	
Target Details		
Target:	CD248	


Target Details

Alternative Name:	CD248 / TEM1 (CD248 Products)	
Background:	Tumor endothelial marker (TEM) 1 was originally identified as a human embryonic fibroblast-	
J	specific antigen and was later determined to be endosialin, a single-pass transmembrane	
	glycoprotein that has multiple extracellular domains, including three EGF-like domains, a sushi-	
	like domain, and a C lectin-like domain. TEM proteins are significantly up-regulated during	
	angiogenesis and neoangiogenesis that are crucial for the growth of solid tumors. While TEM1	
	is not required for angiogenesis during fetal development, postnatal growth or wound healing, it	
	plays a role in tumor growth, invasion, and metastasis. Fibronectin and collagen types I and IV	
	act as specific ligands of TEM1, leading to suggestions that these molecules may cause	
	changes in the extracellular matrix, cell adhesion and migration during tumor invasion. At least	
	two isoforms of TEM1 are known to exist. Synonyms: CD164L1, Endosialin, Tumor endothelial	
	marker 1	
Gene ID:	57124	
NCBI Accession:	NP_065137	
Application Details		
Application Notes:	ELISA. Western blot. Immunohistochemistry on paraffin sections.	
	Other applications not tested.	
	Optimal dilutions are dependent on conditions and should be determined by the user.	
Restrictions:	For Research Use only	
Handling		
Buffer:	PBS, 0.02 % Sodium Azide	
Preservative:	Sodium azide	
Precaution of Use:	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which	
	should be handled by trained staff only.	
Handling Advice:	Avoid repeated freezing and thawing.	
Storage:	4 °C/-20 °C	
Storage Comment:	Store undiluted at 2-8 °C for one month or (in aliquots) at -20 °C for longer.	




Immunofluorescence

Image 1. Immunofluorescence of TEM1 in Human Colon cells with TEM1at 20 μ g/ml.



Immunohistochemistry (Paraffin-embedded Sections)

Image 2. Immunohistochemistry of TEM1 in human colon tissue with this product at $2.5 \, \mu g/ml$.



Western Blotting

Image 3. Western blot analysis of TEM1 in human colon tissue lysate with this product at (A) 0.5 and (B) 1 μ g/ml.