

## Datasheet for ABIN5008861 anti-PISD antibody (AbBy Fluor® 680)



Overview 100 µL Quantity: PISD Target: Reactivity: Human, Mouse, Rat Rabbit Host: Polyclonal Clonality: Conjugate: This PISD antibody is conjugated to AbBy Fluor® 680 Application: Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) **Product Details** KLH conjugated synthetic peptide derived from human PISD Immunogen: Isotype: lgG Cross-Reactivity: Human, Mouse, Rat Purification: Purified by Protein A. Target Details PISD Target: Alternative Name: **PISD (PISD Products)** Background: Synonyms: DJ858B16, dJ858B16.2, DKFZp566G2246, Phosphatidylserine decarboxylase,

Phosphatidylserine decarboxylase beta chain, Phosphatidylserine decarboxylase proenzyme,

PISD, PISD\_HUMAN, PSD, PSDC, PSSC.

Background: Enzymes known as phosphatidylserine decarboxylases (PSDs) catalyze the

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN5008861 | 07/25/2024 | Copyright antibodies-online. All rights reserved.

|                     | formation of phosphatidylethanolamine from phosphatidylserine via phosphatidylserine           |
|---------------------|--|
|                     | decarboxylation. Type I PSDs contain LGST motifs and are found in bacteria and eukaryotic      |
|                     | mitochondria, whereas type II PSDs contain GGST motifs and are found in eukaryotic             |
|                     | endomembrane systems. PISD (phosphatidylserine decarboxylase), also known as                   |
|                     | phosphatidylserine decarboxylase proenzyme, PSDC, PSD, PSSC, DJ858B16, dJ858B16.2 or           |
|                     | DKFZp566G2246, is a 408 amino acid a type I phosphatidylserine decarboxylase that localizes    |
|                     | to the inner mitochondrial membrane. PISD contains a conserved LGST motif which is cleaved     |
|                     | to produce two isoforms known as PISD and PISD $\_$ . PISD is capable of forming a heterodimer |
|                     | and is highly expressed in liver and testis. The gene encoding PISD maps to human              |
|                     | chromosome 22q12.2.  |
| Gene ID:            | 23761  |
| Application Details |  |
| Application Notes:  | IF(IHC-P) 1:50-200   |
| Restrictions:       | For Research Use only  |
| Handling            |  |
| Format:             | Liquid   |
| Concentration:      | 1 µg/µL  |
| Buffer:             | Aqueous buffered solution containing 0.01M TBS ( pH 7.4) with 1 % BSA, 0.03 % Proclin300 and   |
|                     | 50 % Glycerol.   |
| Preservative:       | ProClin  |
| Precaution of Use:  | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be            |
|                     | handled by trained staff only.   |
| Storage:            | -20 °C   |
| Storage Comment:    | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.              |
| Expiry Date:        | 12 months  |

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/2 | Product datasheet for ABIN5008861 | 07/25/2024 | Copyright antibodies-online. All rights reserved.