antibodies .- online.com





anti-PNCK antibody (AA 31-160) (Alexa Fluor 680)



Go to Product page

\sim					
()	VE	۲۱	/1	\triangle	Λ

Quantity:	100 μL
Target:	PNCK
Binding Specificity:	AA 31-160
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PNCK antibody is conjugated to Alexa Fluor 680
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human PNCK
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat,Cow,Sheep,Pig
Purification:	Purified by Protein A.

Target Details

Target:	PNCK
Alternative Name:	PNCK (PNCK Products)
Background:	Synonyms: Calcium/calmodulin dependent protein kinase type 1B, Calcium/calmodulin-

dependent protein kinase type 1B, CaM KI beta, CaM kinase I beta, CaM kinase IB, CaM-KI beta, CaMK1b, CaMKI beta, CaMKI-beta, EC 2.7.11.17, KCC1B_HUMAN, MGC45419, PNCK, Pregnancy up regulated non ubiquitously expressed CaM kinase, Pregnancy up-regulated non-ubiquitously-expressed CaM kinase.

Background: The Ca2+/calmodulin-dependent protein kinases (CaMKs) comprise a structurally related subfamily of serine/threonine kinases. CaMKI Beta (Ca2+/calmodulin-dependent protein kinase type 1B), also known as PNCK (pregnancy up-regulated non-ubiquitously expressed CaM kinase) or BSTK3, is a 343 amino acid protein that localizes to both the nucleus and the cytoplasm and contains one protein kinase domain. Existing as multiple alternatively spliced isoforms, CaMKI Beta functions to catalyze the ATP-dependent phosphorylation of CaMKI, an event that activates CaMKI activity and may be important for Ca2+-triggered signaling cascades within the cell. The gene encoding CaMKI Beta maps to human chromosome X, which contains nearly 153 million base pairs and houses over 1,000 genes.

Application Details

Application Notes:	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	
Expiry Date:	12 months	