# antibodies -online.com





## anti-Prominin 2 antibody (AA 301-400) (Alexa Fluor 680)



Go to Product page

$\sim$			
	N/P	r\/I	i⊢₩

Quantity:	100 μL
Target:	Prominin 2 (PROM2)
Binding Specificity:	AA 301-400
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Prominin 2 antibody is conjugated to Alexa Fluor 680
Application:	Western Blotting (WB), Flow Cytometry (FACS), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human Prominin 2
Isotype:	IgG
Cross-Reactivity:	Human
Predicted Reactivity:	Mouse,Rat,Dog,Pig
Purification:	Purified by Protein A.

#### Target Details

Target:	Prominin 2 (PROM2)
Alternative Name:	Prominin 2 (PROM2 Products)

#### **Target Details**

9	
Background:	Synonyms: PROML2, Prominin-2, PROM-2, Prominin-like protein 2, hPROML2, PROM2,
	UNQ2521/PRO6014
	Background: Prominin 2 is a 112 kDa glycoporotein structurally related to Prominin 1 (CD133)
	although amino acid similarity is not more than 30 $\%$ , but their genomic organization is
	strikingly similar. Like Prominin 1, the prominin 2 exhibit similar membrane topology with 5
	trans-membrane domains and two large glycosylated extracellular domains. Similar to
	Prominin1 localization, the Prominin 2 is also associated with membrane protrusions of the
	epithelial cells from adult kidney, and all along the digestive track and other epithelial
	tissues.Prominin 2 expression is down-regulated in aggressive prostate cancer cell lines and
	transient transfection of PROML2 expression vectors has been shown to induce apoptosis in
	cultured prostate cancer cells, suggesting a tumor suppressive role for Prominin 2. Prominin 2
	expression is likely to be involved in growth suppression in the prostate, and down-regulation of
	Prominin 2 may disrupt normal prostatic homeostasis and lead to uncontrolled prostatic
	growth.
Gene ID:	150696
UniProt:	Q8N271
Application Details	
Application Notes:	FCM 1:20-100
	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only
Llandling	
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
	50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be

handled by trained staff only.

### Handling

Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months