

Datasheet for ABIN5009407

anti-PTPN7 antibody (pSer246) (AbBy Fluor® 680)



Go to Product page

Overview	
Quantity:	100 μL
Target:	PTPN7
Binding Specificity:	pSer246
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PTPN7 antibody is conjugated to AbBy Fluor® 680
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))
Product Details	
Immunogen:	KLH conjugated synthetic phosphopeptide derived from mouse PTPN7 around the phosphorylation site of Ser246
Isotype:	IgG
Cross-Reactivity:	Mouse
Predicted Reactivity:	Rat
Purification:	Purified by Protein A.
Target Details	
Target:	PTPN7

Target Details

Alternative Name:	PTPN7 (PTPN7 Products)
Background:	Synonyms: PTPN7phospho S246, BPTP 4, BPTP4, Dual specicity phosphatase 1,
	Hematopoietic protein tyrosine phosphatase, LC PTP, LCPTP, LPTP, Protein tyrosine
	phoshatase non receptor type stress induced, Protein tyrosine phoshatase nonreceptor type
	stress induced, Protein tyrosine phosphatase LC PTP, Protein tyrosine phosphatase non
	receptor type 7, PTPN 7, PTPNI, Tyrosine protein phosphatase non receptor type 7, PTN7_HUMAN.
	Background: The protein encoded by this gene is a member of the protein tyrosine phosphatase
	(PTP) family. PTPs are known to be signaling molecules that regulate a variety of cellular
	processes including cell growth, differentiation, mitotic cycle, and oncogenic transformation.
	This gene is preferentially expressed in a variety of hematopoietic cells, and is an early
	response gene in lymphokine stimulated cells. The non-catalytic N-terminus of this PTP can
	interact with MAP kinases and suppress the MAP kinase activities. This PTP was shown to be
	involved in the regulation of T cell antigen receptor (TCR) signaling, which was thought to
	function through dephosphorylating the molecules related to MAP kinase pathway. Multiple
	alternatively spliced transcript variants have been found for this gene. [provided by RefSeq].
Gene ID:	5778
Application Details	
Application Notes:	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Drogonyativa	ProClin
Preservative:	

Handling

Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months