

Datasheet for ABIN5009410

anti-PTPN7 antibody (pSer93) (AbBy Fluor® 750)



Go to Product page

Overview		
Quantity:	100 μL	
Target:	PTPN7	
Binding Specificity:	pSer93	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This PTPN7 antibody is conjugated to AbBy Fluor® 750	
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	
Product Details		
Immunogen:	KLH conjugated synthetic phosphopeptide derived from human PTPN7 around the phosphorylation site of Ser93	
Isotype:	IgG	
Predicted Reactivity:	Human,Mouse,Rat,Cow	
Purification:	Purified by Protein A.	
Target Details		
Target:	PTPN7	
Alternative Name:	PTPN7 (PTPN7 Products)	

Target Details

Larget Details		
Background:	Synonyms: PTPN7phospho S93, BPTP 4, BPTP4, Dual specicity phosphatase 1, Hematopoietic	
	protein tyrosine phosphatase, LC PTP, LCPTP, LPTP, Protein tyrosine phoshatase non receptor	
	type stress induced, Protein tyrosine phoshatase nonreceptor type stress induced, Protein	
	tyrosine phosphatase LC PTP, Protein tyrosine phosphatase non receptor type 7, PTPN 7,	
	PTPNI, Tyrosine protein phosphatase non receptor type 7, PTN7_HUMAN.	
	Background: The protein encoded by this gene is a member of the protein tyrosine phosphatase	
	(PTP) family. PTPs are known to be signaling molecules that regulate a variety of cellular	
	processes including cell growth, differentiation, mitotic cycle, and oncogenic transformation.	
	This gene is preferentially expressed in a variety of hematopoietic cells, and is an early	
	response gene in lymphokine stimulated cells. The non-catalytic N-terminus of this PTP can	
	interact with MAP kinases and suppress the MAP kinase activities. This PTP was shown to be	
	involved in the regulation of T cell antigen receptor (TCR) signaling, which was thought to	
	function through dephosphorylating the molecules related to MAP kinase pathway. Multiple	
	alternatively spliced transcript variants have been found for this gene. [provided by RefSeq].	
Gene ID:	5778	
Application Details		
Application Notes:	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and	
	50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be	

Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

handled by trained staff only.

-20 °C

Storage:

Storage Comment:

1	1	l:
-	ココロロ	urnen
1	Hand	III IU

Expiry Date:

12 months