antibodies

## Datasheet for ABIN5009554 anti-GPRC5A antibody (AA 1-33) (AbBy Fluor® 680)



Overview

| Quantity:            | 100 µL   |
|----------------------|--|
| Target:              | GPRC5A   |
| Binding Specificity: | AA 1-33  |
| Reactivity:          | Human  |
| Host:                | Rabbit   |
| Clonality:           | Polyclonal   |
| Conjugate:           | This GPRC5A antibody is conjugated to AbBy Fluor® 680  |
| Application:         | Flow Cytometry (FACS), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |

## Product Details

| Immunogen:        | KLH conjugated synthetic peptide derived from human RAI3 |
|-------------------|--|
| Isotype:          | IgG  |
| Cross-Reactivity: | Human  |
| Purification:     | Purified by Protein A.                                   |

## Target Details

| Target:           | GPRC5A   |
|-------------------|--|
| Alternative Name: | RAI3 (GPRC5A Products)   |
| Background:       | Synonyms: G protein coupled receptor family C group 5 member A, G-protein coupled receptor |

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN5009554 | 06/08/2024 | Copyright antibodies-online. All rights reserved. family C group 5 member A, GPCR 5A, GPCR5A, Gprc 5a, Gprc5a, Orphan G protein coupling receptor PEIG 1, Orphan G protein coupling receptor PEIG1, Orphan G-protein-coupling receptor PEIG-1, RAI 3, RAI3\_HUMAN, Raig 1, RAIG-1, Raig1, Retinoic acid induced 3, Retinoic acid induced gene 1 protein, Retinoic acid induced protein 3, Retinoic acid responsive, Retinoic acid-induced gene 1 protein, Retinoic acid-induced gene 3, Retinoic acid-induced gene 1 protein, Retinoic acid-induced protein 3. Background: RAI3 is an transmembrane G-protein-coupled receptor that affects many essential biological processes including embryogenesis, cell growth, differentiation, and apoptosis. RAI3 may also be involved in maintaining homeostasis of epithelial cells. Retinoic acid receptors directly regulate RAI3 during its transcription in embryonal carcinoma differentiation. RAI3 expression is upregulated in most tumor cell lines that express mutant p53, suggesting that p53 interacts with the promoter of RAI3 and represses its expression at the beginning of apoptosis. RAI3 is a potential molecular target for diagnosing breast cancer, and selective suppression of signals from RAI3 may have a place in breast cancer treatments.

## Application Details

| Application Notes: | FCM 1:20-100   |
|--------------------|--|
|                    | IF(IHC-P) 1:50-200   |
|                    | IF(IHC-F) 1:50-200   |
|                    | IF(ICC) 1:50-200   |
| Restrictions:      | For Research Use only  |
| Handling           |  |
| Format:            | Liquid   |
| Concentration:     | 1 μg/μL  |
| Buffer:            | Aqueous buffered solution containing 0.01M TBS ( pH 7.4) with 1 % BSA, 0.03 % Proclin300 and |
|                    | 50 % Glycerol.   |
| Preservative:      | ProClin  |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be          |
|                    | handled by trained staff only.   |
| Storage:           | -20 °C   |
| Storage Comment:   | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.            |
| Expiry Date:       | 12 months  |

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/2 | Product datasheet for ABIN5009554 | 06/08/2024 | Copyright antibodies-online. All rights reserved.