



[Go to Product page](#)

Datasheet for ABIN5009592

## anti-RAPTOR antibody (pSer792) (AbBy Fluor® 680)

### Overview

Quantity:	100 µL
Target:	RAPTOR
Binding Specificity:	pSer792
Reactivity:	Human, Mouse, Zebrafish (Danio rerio)
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This RAPTOR antibody is conjugated to AbBy Fluor® 680
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

### Product Details

Immunogen:	KLH conjugated synthetic phosphopeptide derived from mouse Raptor around the phosphorylation site of Ser792
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Zebrafish (Danio rerio)
Predicted Reactivity:	Rat,Dog,Cow,Sheep,Pig,Horse,Chicken,Guinea Pig
Purification:	Purified by Protein A.

### Target Details

Target:	RAPTOR
---------	--------

## Target Details

---

Alternative Name: Raptor ([RAPTOR Products](#))

---

Background: Synonyms: ARaptor phospho S792, Raptor phospho Ser792, p-Raptor Ser792, p150 target of rapamycin TOR scaffold protein containing WD repeats, P150 target of rapamycin TOR-scaffold protein, Regulatory Associated Protein of mTOR, KIAA1303, RPTOR\_HUMAN, Regulatory-associated protein of mTOR, p150 target of rapamycin TOR-scaffold protein.

Background: mTOR controls cell growth, in part by regulating p70 S6 kinase alpha (p70alpha) and eukaryotic initiation factor 4E binding protein 1 (4EBP1). Raptor is a 150 kDa mTOR binding protein that also binds 4EBP1 and p70alpha. The binding of Raptor to mTOR is necessary for the mTOR-catalyzed phosphorylation of 4EBP1 in vitro, and it strongly enhances the mTOR kinase activity toward p70alpha. Rapamycin or amino acid withdrawal increases, whereas insulin strongly inhibits, the recovery of 4EBP1 and raptor on 7-methyl-GTP Sepharose. Partial inhibition of raptor expression by RNA interference (RNAi) reduces mTOR-catalyzed 4EBP1 phosphorylation in vitro. RNAi of *C. elegans* raptor yields an array of phenotypes that closely resemble those produced by inactivation of Ce-TOR. Thus, raptor is an essential scaffold for the mTOR-catalyzed phosphorylation of 4EBP1 and mediates TOR action in vivo.

---

Pathways: [Warburg Effect](#)

---

## Application Details

---

Application Notes: IF(IHC-P) 1:50-200  
IF(IHC-F) 1:50-200  
IF(ICC) 1:50-200

---

Restrictions: For Research Use only

---

## Handling

---

Format: Liquid

---

Concentration: 1 µg/µL

---

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

---

Preservative: ProClin

---

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

---

Storage: -20 °C

---

## Handling

---

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

---

Expiry Date: 12 months