## antibodies .- online.com





## anti-REC8 antibody (pSer251) (Alexa Fluor 680)



Go to Product page

| Overview             |  |
|----------------------|--|
| Quantity:            | 100 μL   |
| Target:              | REC8   |
| Binding Specificity: | pSer251  |
| Reactivity:          | Mouse, Rat   |
| Host:                | Rabbit   |
| Clonality:           | Polyclonal   |
| Conjugate:           | This REC8 antibody is conjugated to Alexa Fluor 680  |
| Application:         | Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |
| Product Details      |  |
| Immunogen:           | KLH conjugated synthetic phosphopeptide derived from mouse REC8 around the phosphorylation site of Ser251                      |
| Isotype:             | IgG  |
| Cross-Reactivity:    | Mouse, Rat   |
| Purification:        | Purified by Protein A.   |
| Target Details       |  |
| Target:              | REC8   |
| Alternative Name:    | REC8 (REC8 Products)   |

## **Target Details**

| raiget betails      |   |  |
|---------------------|---|--|
| Background:         | Synonyms: mrec, Rec8L1, Meiotic recombination protein REC8 homolog, Cohesin Rec8p, Rec8, Mei8 |  |
|                     | Background: Required during meiosis for separation of sister chromatids and homologous        |  |
|                     | chromosomes. Proteolytic cleavage of REC8 on chromosome arms by separin during                |  |
|                     | anaphase I allows for homologous chromosome separation in meiosis I and cleavage of REC8      |  |
|                     | on centromeres during anaphase II allows for sister chromatid separation in meiosis II.       |  |
| Gene ID:            | 56739   |  |
| UniProt:            | Q8C5S7  |  |
| Application Details |   |  |
| Application Notes:  | IF(IHC-P) 1:50-200  |  |
|                     | IF(IHC-F) 1:50-200  |  |
|                     | IF(ICC) 1:50-200  |  |
| Restrictions:       | For Research Use only   |  |
| Handling            |   |  |
| Format:             | Liquid  |  |
| Concentration:      | 1 μg/μL   |  |
| Buffer:             | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and   |  |
|                     | 50 % Glycerol.  |  |
| Preservative:       | ProClin   |  |
| Precaution of Use:  | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be           |  |
|                     | handled by trained staff only.  |  |
| Storage:            | -20 °C  |  |
| Storage Comment:    | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.             |  |
| Expiry Date:        | 12 months   |  |