antibodies.com

## Datasheet for ABIN5009912 anti-SH3RF1 antibody (Alexa Fluor 750)



Overview	
Quantity:	100 μL
Target:	SH3RF1
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This SH3RF1 antibody is conjugated to Alexa Fluor 750
Application:	Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p))
Product Details	
Immunogen:	KLH conjugated synthetic peptide derived from human SH3MD2/RNF142
lsotype:	lgG
Cross-Reactivity:	Human, Mouse, Rat
Purification:	Purified by Protein A.
Target Details	
Target:	SH3RF1
Alternative Name:	RNF142 (SH3RF1 Products)
Background:	Synonyms: Plenty of SH3 domains, Plenty of SH3s, POSH, Ring finger protein 142, RNF 142,
	RNF142, SH3 domain containing ring finger 1, SH3 domain containing RING finger protein 1,
	SH3MD2, SH3 multiple domains 2, SH3RF1.
	Background: SH3MD2 acts as a scaffold protein, contributes to Rac-induced signal

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN5009912 | 03/07/2024 | Copyright antibodies-online. All rights reserved.

	transduction such as JNKs (MAPK8 and MAPK9) activation and induces apoptosis. Within a
	signaling complex, it probably recruits protein kinases such as MAP3K10 or MAP3K11 which
	are in turn activated leading to the sequential activation of MAP2K4, MAP2K7 and JNKs
	(MAPK8 and MAPK9). SH3MD2 may be involved in targeting of HIV-1 GAG and GAG-POL
	polyproteins to the plasma membrane. This gene encodes a protein containing an N-terminus
	RING-finger, four SH3 domains, and a region implicated in binding of the Rho GTPase Rac. Via
	the RING-finger, the encoded protein has been shown to function as an ubiquitin-protein ligase
	involved in protein sorting at the trans-Golgi network. The encoded protein may also act as a
	scaffold for the c-Jun N-terminal kinase signaling pathway, facilitating the formation of a
	functional signaling module. There are two named isoforms.
Gene ID:	57630
Application Details	
Application Notes:	IF(IHC-P) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS ( pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
	50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
	handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months