# antibodies -online.com





## anti-DLGAP2 antibody (AA 251-360) (AbBy Fluor® 750)



Go to Product page

$\sim$			
	N/P	r\/	i⊢₩

Quantity:	100 μL
Target:	DLGAP2
Binding Specificity:	AA 251-360
Reactivity:	Human, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This DLGAP2 antibody is conjugated to AbBy Fluor® 750
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

## **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human SAPAP2/DLGAP2
Isotype:	IgG
Cross-Reactivity:	Human, Rat
Predicted Reactivity:	Mouse
Purification:	Purified by Protein A.

## **Target Details**

Target:	DLGAP2
Alternative Name:	SAPAP2/DLGAP2 (DLGAP2 Products)

#### **Target Details**

#### Background:

Synonyms: DAP-2, DAP2, Discs large Drosophila homolog associated protein 2, Discs large homolog associated protein 2, Disks large-associated protein 2, Dlgap2, DLGP2\_HUMAN, PSD-95/SAP90-binding protein 2, SAP90/PSD-95-associated protein 2, SAPAP2, hide. Background: A guanylate kinase is a phosphotransferase that produces ADP and GDP from the substrates ATP and GMP. SAPAP2, also known as DAP-2 (Disks large-associated protein 2) and PSD-95/SAP90-binding protein 2, is a 1054 amino acid protein that localizes to the postsynaptic membrane of neuronal cells of the brain and kidney. SAPAP2 likely acts as a signaling molecule which interacts with the human genes DLG1 and DLG4/PSD-95. The gene encoding SAPAP2, DLGAP2, maps to human chromosome 8. Consisting of nearly 146 million base pairs, chromosome 8 encodes over 800 genes and is associated with a variety of diseases and malignancies. Schizophrenia, bipolar disorder, Trisomy 8, Pfeiffer syndrome, congenital hypothyroidism, Waardenburg syndrome and some leukemias and lymphomas are thought to occur as a result of defects in specific genes that maps to chromosome 8.

### **Application Details**

Application Notes:	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	
Expiry Date:	12 months	