-online.com antibodies

## Datasheet for ABIN5010244 anti-SASS6 antibody (Alexa Fluor 680)

μL



Quantity:	100
Target:	SAS

Overview

Target:	SASS6
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This SASS6 antibody is conjugated to Alexa Fluor 680
Application:	Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

## Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human SASS6
lsotype:	lgG
Cross-Reactivity:	Human, Mouse, Rat
Purification:	Purified by Protein A.

## Target Details

Target:	SASS6
Alternative Name:	SASS6 (SASS6 Products)
Background:	Synonyms: HsSAS 6, HsSAS6, MGC119440, SAS 6, SAS6, SASS 6, Spindle assembly 6 homolog
	C. elegans, Spindle assembly 6 homolog, Spindle assembly abnormal protein 6 homolog,
	PJA2_HUMAN.
	Background: SAS-6 (spindle assembly abnormal protein 6 homolog, HsSAS-6) is a 657 amino

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN5010244 | 03/07/2024 | Copyright antibodies-online. All rights reserved.

	acid protein encoded by the human gene SAS6. SAS-6 is a component of the centrosome that
	contains one PISA (present in SAS-6) domain. LK4, SAS-6, CPAP and other centriole related
	proteins are required at different stages of procentriole formation and were associated with
	different centriolar structures. SAS-6 associates only transiently with nascent procentrioles,
	whereas CEP135 and CPAP form a core structure within the proximal lumen of both parental
	and nascent centrioles. SAS-6 is necessary for procentriole formation in human cell lines and is
	localized asymmetrically next to the centriole at the onset of procentriole formation. SAS-6
	levels oscillate during the cell cycle, it is degraded in mitosis starting at anaphase, and it
	accumulates again at the end of the following G1 phase. The anaphase-promoting complex
	targets SAS-6 for degradation by the 26S Proteasome, and a KEN box in the C-terminus of SAS-
	6 is necessary for its degradation. Increased SAS-6 levels promoted the formation of multiple
	procentrioles forming next to a single centriole.
Gene ID:	163786

## Application Details

Application Notes:	IF(IHC-P) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid

Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS ( pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/2 | Product datasheet for ABIN5010244 | 03/07/2024 | Copyright antibodies-online. All rights reserved.