antibodies -online.com





anti-SEMA6D antibody (AA 801-900) (Alexa Fluor 680)



Go to Product page

| () | 1/0 | r\ /1 | 014 | |
|-----|-----|-------|-----|---|
| () | ve | I V I | -v | V |
| | | | | |

| Quantity: | 100 μL | |
|----------------------|---|--|
| Target: | SEMA6D | |
| Binding Specificity: | AA 801-900 | |
| Reactivity: | Mouse, Rat | |
| Host: | Rabbit | |
| Clonality: | Polyclonal | |
| Conjugate: | This SEMA6D antibody is conjugated to Alexa Fluor 680 | |
| Application: | Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) | |

Product Details

| Immunogen: | KLH conjugated synthetic peptide derived from human SEMA6D | |
|-----------------------|--|--|
| Isotype: | IgG | |
| Cross-Reactivity: | Mouse, Rat | |
| Predicted Reactivity: | Human,Dog,Cow,Pig,Horse | |
| Purification: | Purified by Protein A. | |

Target Details

| Target: | SEMA6D |
|-------------------|--------------------------|
| Alternative Name: | SEMA6D (SEMA6D Products) |

Target Details

| De alcemano: di | Oursen was OFMCD HIIIMAN KIAA4470 OONEVA Ooroo da | | |
|---------------------|---|--|--|
| Background: | Synonyms: SEM6D_HUMAN, KIAA1479, Q8NFY4, Sema domain, transmembrane domain TM, | | |
| | and cytoplasmic domain, semaphorin 6D, Semaphorin 6D. | | |
| | Background: Semaphorins are a large family, including both secreted and membrane | | |
| | associated proteins, many of which have been implicated as inhibitors or chemorepellents in | | |
| | axon pathfinding, fasciculation and branching, and target selection. All semaphorins possess a | | |
| | semaphorin (Sema) domain and a PSI domain (found in plexins, semaphorins and integrins) in | | |
| | the N-terminal extracellular portion. Additional sequence motifs C-terminal to the semaphorin | | |
| | domain allow classification into distinct subfamilies. Results demonstrate that transmembrane | | |
| | semaphorins, like the secreted ones, can act as repulsive axon guidance cues. This gene | | |
| | encodes a class 6 vertebrate transmembrane semaphorin that demonstrates alternative | | |
| | splicing. Several transcript variants have been identified and expression of the distinct encoded | | |
| | isoforms is thought to be regulated in a tissue- and development-dependent manner. [provided | | |
| | by RefSeq, Nov 2010] | | |
| Gene ID: | 80031 | | |
| Pathways: | Smooth Muscle Cell Migration | | |
| Application Details | | | |
| Application Notes: | IF(IHC-P) 1:50-200 | | |
| | IF(IHC-F) 1:50-200 | | |
| | IF(ICC) 1:50-200 | | |
| Restrictions: | For Research Use only | | |
| Handling | | | |
| Format: | Liquid | | |
| Concentration: | 1 μg/μL | | |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and | | |
| | 50 % Glycerol. | | |
| Preservative: | ProClin | | |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be | | |
| | handled by trained staff only. | | |
| Storage: | -20 °C | | |
| Storage Comment: | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles. | | |
| | Store at 20 0.7 inquot into martiple vialo to avoid repeated neeze than oyoleo. | | |

| Н | land | lına |
|---|-------|------|
| | iaria | шц |

Expiry Date:

12 months