

Datasheet for ABIN5010615

anti-SLC25A11 antibody (AA 35-140) (AbBy Fluor® 750)



Go to Product page

()	V		rV	ĺ	9	V	V
'	\mathcal{I}	٧V	<u> </u>	v	1	$\overline{}$	٧	٧

Target: SLC25A11 Binding Specificity: AA 35-140 Reactivity: Human Host: Rabbit Clonality: Polyclonal Conjugate: This SLC25A11 antibody is conjugated to AbBy Fluor® 750 Application: Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded	Quantity:	100 μL	
Reactivity: Human Host: Rabbit Clonality: Polyclonal Conjugate: This SLC25A11 antibody is conjugated to AbBy Fluor® 750 Application: Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded	Target:	SLC25A11	
Host: Rabbit Clonality: Polyclonal Conjugate: This SLC25A11 antibody is conjugated to AbBy Fluor® 750 Application: Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded	Binding Specificity:	AA 35-140	
Clonality: Polyclonal Conjugate: This SLC25A11 antibody is conjugated to AbBy Fluor® 750 Application: Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded	Reactivity:	Human	
Conjugate: This SLC25A11 antibody is conjugated to AbBy Fluor® 750 Application: Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded	Host:	Rabbit	
Application: Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded	Clonality:	Polyclonal	
	Conjugate:	This SLC25A11 antibody is conjugated to AbBy Fluor® 750	
	Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded	
Sections) (IF (p))		Sections) (IF (p))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human SLC25A11		
Isotype:	IgG		
Cross-Reactivity:	Human		
Predicted Reactivity:	Mouse,Rat,Dog,Cow,Pig,Horse,Rabbit		
Purification:	Purified by Protein A.		

Target Details

Target:	SLC25A11
Alternative Name:	SLC25A11 (SLC25A11 Products)

Target Details

Synonyms: Mitochondrial 2 oxoglutarate/malate carrier protein, OGC, OGCP, SLC20A4, Solute
carrier family 20 oxoglutarate carrier member 4, Solute carrier family 20 member 4, Solute
carrier family 25 mitochondrial carrier oxoglutarate carrier member 11, Solute carrier family 25
member 11.
Background: SLC25A11 catalyzes the transport of 2-oxoglutarate across the inner
mitochondrial membrane in an electroneutral exchange for malate or other dicarboxylic acids,
and plays an important role in several metabolic processes, including the malate-aspartate
shuttle, the oxoglutarate/isocitrate shuttle, in gluconeogenesis from lactate, and in nitrogen
metabolism.
8402
Dicarboxylic Acid Transport
IF(IHC-P) 1:50-200
IF(IHC-F) 1:50-200
IF(ICC) 1:50-200
For Research Use only
Liquid
1 μg/μL
Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
50 % Glycerol.
ProClin
This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
handled by trained staff only.
-20 °C
Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
12 months