# antibodies -online.com





# anti-SLC37A4 antibody (AA 25-130) (Alexa Fluor 680)



Go to Product page

| $\sim$ |     |     |     |
|--------|-----|-----|-----|
|        | N/P | r\/ | i⊢₩ |

| Quantity:            | 100 μL   |  |
|----------------------|--|--|
| Target:              | SLC37A4  |  |
| Binding Specificity: | AA 25-130  |  |
| Reactivity:          | Human  |  |
| Host:                | Rabbit   |  |
| Clonality:           | Polyclonal   |  |
| Conjugate:           | This SLC37A4 antibody is conjugated to Alexa Fluor 680   |  |
| Application:         | Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |  |

## Product Details

| Immunogen:            | KLH conjugated synthetic peptide derived from human SLC37A4 |
|-----------------------|---|
| Isotype:              | IgG   |
| Cross-Reactivity:     | Human   |
| Predicted Reactivity: | Mouse,Rat,Dog,Cow,Pig                                       |
| Purification:         | Purified by Protein A.                                      |

### **Target Details**

| Target:           | SLC37A4                    |
|-------------------|----------------------------|
| Alternative Name: | SLC37A4 (SLC37A4 Products) |

### **Target Details**

| Synonyms: G6PT2, GSD1b, GSD1c, GSD1d, TRG19, G6PT1, G6PT3, Glucose-5-phosphate                 |  |
|--|--|
|  |  |
| transporter, Glucose-6-phosphatase, transport glucose protein 3 antibody, Glucose-6-           |  |
| phosphatase, transport glucose-6-phosphate protein 1, Glucose-6-phosphatase, transport         |  |
| phosphate/pyrophosphate protein 2, Glucose-6-phosphate translocase, Glucose-6-phosphate        |  |
| transporter 1, Microsomal glucose-6-phosphate transporter, Solute carrier family 37 glucose-6- |  |
| phosphate transporter, member 4, MGC15729, PRO0685, G6PT1_HUMAN.                               |  |
| Background: SLC37A4 transports glucose-6-phosphate from the cytoplasm to the lumen of the      |  |
| endoplasmic reticulum. It forms a complex with glucose-6-phosphatase which is responsible      |  |
| for glucose production through glycogenolysis and gluconeogenesis. Hence, it plays a central   |  |
| role in homeostatic regulation of blood glucose levels.  |  |
| 2542   |  |
| Carbohydrate Homeostasis, Cellular Glucan Metabolic Process                                    |  |
|  |  |
| IF(IHC-P) 1:50-200   |  |
| IF(IHC-F) 1:50-200   |  |
| IF(ICC) 1:50-200   |  |
| For Research Use only  |  |
|  |  |
| Liquid   |  |
| 1 μg/μL  |  |
| Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and    |  |
| 50 % Glycerol.   |  |
| ProClin  |  |
| This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be            |  |
| handled by trained staff only.   |  |
| -20 °C   |  |
| Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.              |  |
| 12 months  |  |
|  |  |