



[Go to Product page](#)

Datasheet for ABIN5010645

anti-SLC38A2 antibody (AA 21-150) (Alexa Fluor 750)

Overview

Quantity:	100 µL
Target:	SLC38A2
Binding Specificity:	AA 21-150
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This SLC38A2 antibody is conjugated to Alexa Fluor 750
Application:	Western Blotting (WB), Flow Cytometry (FACS), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human SLC38A2/SNAT2
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Predicted Reactivity:	Dog,Cow,Sheep,Pig,Horse
Purification:	Purified by Protein A.

Target Details

Target:	SLC38A2
Alternative Name:	SLC38A2/SNAT2 (SLC38A2 Products)

Target Details

Background: Synonyms: Amino acid transporter 2, Amino acid transporter A2, ATA2, KIAA1382, PRO1068, Protein 40-9-1, S38A2_HUMAN, SAT2, Slc38a2, SNAT2, Sodium-coupled neutral amino acid transporter 2, Solute carrier family 38 member 2, System A amino acid transporter, System A amino acid transporter 2, System A transporter 1, System N amino acid transporter 2.

Background: The sodium-coupled neutral amino acid transporters (SNAT) of the SLC38 gene family include System A subtypes SNAT1, SNAT2 and SNAT4 and System N subtypes SNAT3 and SNAT5. The SLC38 transporters are essential for the uptake of nutrients, energy production, metabolism, detoxification, and the cycling of neurotransmitters. SNAT2, also designated ATA2, PRO1068 and SAT2 is encoded by the human gene SLC38A2. The functional role of SNAT2 in the nervous system is unclear. Protein expression is notably enriched in the spinal cord and brain stem nuclei of the auditory system. System A transport proteins are also present in placental tissue. These SNAT proteins may play a significant role in fetal development and inhibition of the transport system has been associated with fetal growth retardation.

UniProt: [Q96QD8](#)

Pathways: [Dicarboxylic Acid Transport](#)

Application Details

Application Notes: FCM 1:20-100
IF(IHC-P) 1:50-200
IF(IHC-F) 1:50-200
IF(ICC) 1:50-200

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Handling

Storage: -20 °C

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Expiry Date: 12 months