# antibodies -online.com





# anti-SNAPIN antibody (AA 51-136) (Alexa Fluor 750)



Go to Product page

| $\sim$ |          |
|--------|----------|
| ( )\/△ | rview    |
| $\cup$ | 1 410 44 |

| Quantity:            | 100 μL  |
|----------------------|---|
| Target:              | SNAPIN  |
| Binding Specificity: | AA 51-136   |
| Reactivity:          | Human   |
| Host:                | Rabbit  |
| Clonality:           | Polyclonal  |
| Conjugate:           | This SNAPIN antibody is conjugated to Alexa Fluor 750   |
| Application:         | Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |

### **Product Details**

| Immunogen:            | KLH conjugated synthetic peptide derived from human SNAP25BP |
|-----------------------|--|
| Isotype:              | IgG  |
| Predicted Reactivity: | Human, Mouse, Rat, Dog, Cow, Pig, Horse, Chicken, Rabbit     |
| Purification:         | Purified by Protein A.                                       |

# **Target Details**

| Target:           | SNAPIN   |
|-------------------|--|
| Alternative Name: | SNAPIN/SNAP25BP (SNAPIN Products)  |
| Background:       | Synonyms: SNAP associated protein, SNAP-25-binding protein, SNAP-associated protein, |

| SNAP25BP, SNAPAP, SNAPIN, SNAPN_HUMAN, SNARE associated protein snapin, SNARE-               |
|--|
| associated protein Snapin, Synaptosomal associated protein 25 binding protein, Synaptosomal- |
| associated protein 25-binding protein, OTTHUMP00000035157.                                   |

Background: The protein encoded by this gene is a coiled-coil-forming protein that associates with the SNARE (soluble N-ethylmaleimide-sensitive fusion protein attachment protein receptor) complex of proteins and the BLOC-1 (biogenesis of lysosome-related organelles) complex. Biochemical studies have identified additional binding partners. As part of the SNARE complex, it is required for vesicle docking and fusion and regulates neurotransmitter release. The BLOC-1 complex is required for the biogenesis of specialized organelles such as melanosomes and platelet dense granules. Mutations in gene products that form the BLOC-1 complex have been identified in mouse strains that are models of Hermansky-Pudlak syndrome. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jun 2012].

Gene ID: 23557

Pathways: Synaptic Membrane, Synaptic Vesicle Exocytosis

### **Application Details**

| Application Notes: | IF(IHC-P) 1:50-200 |
|--------------------|--------------------|
|                    |                    |

IF(IHC-F) 1:50-200

IF(ICC) 1:50-200

Restrictions: For Research Use only

## Handling

| Format:            | Liquid   |
|--------------------|--|
| Concentration:     | 1 μg/μL  |
| Buffer:            | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.         |
| Preservative:      | ProClin  |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. |
| Storage:           | -20 °C   |
| Storage Comment:   | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.                                  |

|          |       | 1.   |   |
|----------|-------|------|---|
| $\vdash$ | land  | lın  |   |
|          | ıaııu | 1111 | U |
|          |       |      |   |

Expiry Date:

12 months