antibodies -online.com







anti-SNIP1 antibody (AA 301-396) (AbBy Fluor® 680)



()	ve	K\ /		
	\cup	1 V/	Щ.	V۷

Quantity:	100 μL	
Target:	SNIP1	
Binding Specificity:	AA 301-396	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This SNIP1 antibody is conjugated to AbBy Fluor® 680	
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human SNIP1
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat,Dog,Cow,Sheep,Pig,Chicken,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	SNIP1
Alternative Name:	SNIP1 (SNIP1 Products)
Background: Synonyms: FHA domain-containing protein SNIP1, FLJ12553, Smad nuclear intera	

Smad nuclear interacting, Smad nuclear interacting protein, Smad nuclear-interacting protein 1,
SNIP1 Smad nuclear interacting protein, SNIP1, SNIP1_HUMAN, Splicing factor arginine/serine
rich 4 Pre mRNA splicing factor SRP75.

Background: Members of the transforming growth factor-beta (TGF-Beta) superfamily play critical roles in controlling cell growth and differentiation. Effects of TGF-Beta family ligands are mediated by Smad proteins. The Smad nuclear interacting protein (SNIP1) contains a forkhead-associated (FHA) domain and acts as a nuclear inhibitor of CBP/p300. SNIP1 potently inhibits the activity of NF-kappa B, which binds the C/H1 domain of CBP/p300, by competing for the binding site. SNIP1 is also thought to induce expression of Cyclin D1 to promote cellular proliferation. SNIP1 is ubiquitously expressed with high expression in heart and skeletal muscle.

Pathways:

SARS-CoV-2 Protein Interactome, The Global Phosphorylation Landscape of SARS-CoV-2 Infection

Application Details

Restrictions:	For Research Use only
	IF(ICC) 1:50-200
	IF(IHC-F) 1:50-200
Application Notes:	IF(IHC-P) 1:50-200

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months