

Datasheet for ABIN5010866

anti-SOX30 antibody (AA 351-450) (AbBy Fluor® 750)



[Go to Product page](#)

Overview

| | |
|----------------------|--|
| Quantity: | 100 µL |
| Target: | SOX30 |
| Binding Specificity: | AA 351-450 |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This SOX30 antibody is conjugated to AbBy Fluor® 750 |
| Application: | Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |

Product Details

| | |
|-----------------------|---|
| Immunogen: | KLH conjugated synthetic peptide derived from human SOX30 |
| Isotype: | IgG |
| Predicted Reactivity: | Human, Mouse, Rat, Dog, Cow, Sheep, Pig |
| Purification: | Purified by Protein A. |

Target Details

| | |
|-------------------|---|
| Target: | SOX30 |
| Alternative Name: | SOX30 (SOX30 Products) |
| Background: | Synonyms: Sex determining region Y box 30, Sox30 protein type II, SRY box 30, SRY related |

Target Details

HMG box 30, Transcription factor SOX 30.

Background: Sox-30 encodes a member of the Sox (SRY-related HMG-box) family of transcription factors involved in the regulation of embryonic development and in the determination of the cell fate. Sox-30 is expressed in testis and may act as a transcriptional regulator after forming a protein complex with other proteins. Sox-30 may be involved in the differentiation of developing male germ cells. Two transcript variants encoding distinct isoforms have been identified for the human Sox-30 gene. Sox family transcription factors influence cell differentiation, development and sex determination. Sox-30 contains a unique DNA binding domain, known as the high mobility group (HMG) box, that is related to that of the testis determining gene, SRY. The highly complex group of Sox genes cluster at a minimum of 40 different loci that rapidly diverged in various animal lineages. Several Sox genes have been identified, and members of this family have been shown to be conserved during evolution and to play key roles during animal development.

Application Details

Application Notes: IF(IHC-P) 1:50-200
IF(IHC-F) 1:50-200
IF(ICC) 1:50-200

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Expiry Date: 12 months