# antibodies .- online.com





# anti-PPP1R9B antibody (AA 358-460) (Alexa Fluor 680)



| ( ) | 1 / | $\sim$   | KI / | 110 | Νę    |
|-----|-----|----------|------|-----|-------|
|     | 1// | $\vdash$ | I \/ | 1 ← | ٠// ٢ |
|     |     |          |      |     |       |

| Quantity:            | 100 μL   |
|----------------------|--|
| Target:              | PPP1R9B  |
| Binding Specificity: | AA 358-460   |
| Reactivity:          | Rat, Mouse   |
| Host:                | Rabbit   |
| Clonality:           | Polyclonal   |
| Conjugate:           | This PPP1R9B antibody is conjugated to Alexa Fluor 680   |
| Application:         | Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |

#### **Product Details**

| Immunogen:            | KLH conjugated synthetic peptide derived from human Spinophilin/Neurabin 2 |
|-----------------------|--|
| Isotype:              | IgG  |
| Cross-Reactivity:     | Mouse, Rat   |
| Predicted Reactivity: | Human,Dog,Sheep,Pig  |
| Purification:         | Purified by Protein A.   |

## **Target Details**

| Target:           | PPP1R9B                                   |
|-------------------|---|
| Alternative Name: | Spinophilin/Neurabin 2 (PPP1R9B Products) |

#### **Target Details**

| Background: |
|-------------|
|-------------|

Synonyms: Spn, SPINO, PPP1R6, PPP1R9, Neurabin-2, Neurabin-II, Protein phosphatase 1 regulatory subunit 9B, Spinophilin, PPP1R9B

Background: Seems to act as a scaffold protein in multiple signaling pathways. Modulates excitatory synaptic transmission and dendritic spine morphology. Binds to actin filaments (F-actin) and shows cross-linking activity. Binds along the sides of the F-actin. May play an important role in linking the actin cytoskeleton to the plasma membrane at the synaptic junction. Believed to target protein phosphatase 1/PP1 to dendritic spines, which are rich in F-actin, and regulates its specificity toward ion channels and other substrates, such as AMPA-type and NMDA-type glutamate receptors. Plays a role in regulation of G-protein coupled receptor signaling, including dopamine D2 receptors and alpha-adrenergic receptors. May establish a signaling complex for dopaminergic neurotransmission through D2 receptors by linking receptors downstream signaling molecules and the actin cytoskeleton. Binds to ADRA1B and RGS2 and mediates regulation of ADRA1B signaling. May confer to Rac signaling specificity by binding to both, RacGEFs and Rac effector proteins. Probably regulates p70 S6 kinase activity by forming a complex with TIAM1 (By similarity). Required for hepatocyte growth factor (HGF)-induced cell migration.

| Gene ID: | 84687  |
|----------|--------|
| UniProt: | Q96SB3 |

Pathways:

Regulation of G-Protein Coupled Receptor Protein Signaling

#### **Application Details**

Application Notes: IF(IHC-P) 1:50-200
IF(IHC-F) 1:50-200
IF(ICC) 1:50-200

Restrictions: Fe

For Research Use only

#### Handling

| Format:        | Liquid   |
|----------------|--|
| Concentration: | 1 μg/μL  |
| Buffer:        | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol. |
| Preservative:  | ProClin  |

## Handling

| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. |
|--------------------|--|
| Storage:           | -20 °C   |
| Storage Comment:   | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.                                  |
| Expiry Date:       | 12 months  |