antibodies .- online.com





anti-PPP1R9B antibody (AA 358-460) (Alexa Fluor 750)



Go to Product page

\sim			
	N/P	r\/	i⊢₩

Quantity:	100 μL	
Target:	PPP1R9B	
Binding Specificity:	AA 358-460	
Reactivity:	Rat, Mouse	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This PPP1R9B antibody is conjugated to Alexa Fluor 750	
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human Spinophilin/Neurabin 2	
Isotype:	IgG	
Cross-Reactivity:	Mouse, Rat	
Predicted Reactivity:	Human,Dog,Sheep,Pig	
Purification:	Purified by Protein A.	

Target Details

Target:	PPP1R9B	
Alternative Name:	Spinophilin/Neurabin 2 (PPP1R9B Products)	

Target Details

Back	aroi	ınd:
Dack	yı Ot	ii iu.

Synonyms: Spn, SPINO, PPP1R6, PPP1R9, Neurabin-2, Neurabin-II, Protein phosphatase 1 regulatory subunit 9B, Spinophilin, PPP1R9B

Background: Seems to act as a scaffold protein in multiple signaling pathways. Modulates excitatory synaptic transmission and dendritic spine morphology. Binds to actin filaments (F-actin) and shows cross-linking activity. Binds along the sides of the F-actin. May play an important role in linking the actin cytoskeleton to the plasma membrane at the synaptic junction. Believed to target protein phosphatase 1/PP1 to dendritic spines, which are rich in F-actin, and regulates its specificity toward ion channels and other substrates, such as AMPA-type and NMDA-type glutamate receptors. Plays a role in regulation of G-protein coupled receptor signaling, including dopamine D2 receptors and alpha-adrenergic receptors. May establish a signaling complex for dopaminergic neurotransmission through D2 receptors by linking receptors downstream signaling molecules and the actin cytoskeleton. Binds to ADRA1B and RGS2 and mediates regulation of ADRA1B signaling. May confer to Rac signaling specificity by binding to both, RacGEFs and Rac effector proteins. Probably regulates p70 S6 kinase activity by forming a complex with TIAM1 (By similarity). Required for hepatocyte growth factor (HGF)-induced cell migration.

Gene ID:

84687

UniProt:

Q96SB3

Pathways:

Regulation of G-Protein Coupled Receptor Protein Signaling

Application Details

Application Notes:

IF(IHC-P) 1:50-200

IF(IHC-F) 1:50-200

IF(ICC) 1:50-200

Restrictions:

For Research Use only

Handling

Format:

Liquid

Concentration:

 $1 \mu g/\mu L$

Buffer:

Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and

50 % Glycerol.

Preservative:

ProClin

Handling

Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months