antibodies - online.com







anti-SRPX2 antibody (AA 121-220) (AbBy Fluor® 750)



| \sim | | | |
|--------|-----|------|-----|
| | N/P | r\/I | i⊢₩ |

| Quantity: | 100 μL | |
|----------------------|--|--|
| Target: | SRPX2 | |
| Binding Specificity: | AA 121-220 | |
| Reactivity: | Mouse | |
| Host: | Rabbit | |
| Clonality: | Polyclonal | |
| Conjugate: | This SRPX2 antibody is conjugated to AbBy Fluor® 750 | |
| Application: | Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) | |

Product Details

| Immunogen: | KLH conjugated synthetic peptide derived from human SRPX2 | |
|-----------------------|---|--|
| Isotype: | IgG | |
| | | |
| Cross-Reactivity: | Mouse | |
| Predicted Reactivity: | Human,Rat | |
| Purification: | Purified by Protein A. | |

Target Details

| Target: | SRPX2 |
|-------------------|------------------------|
| Alternative Name: | SRPX2 (SRPX2 Products) |

Target Details

| Bacl | / 0 | ro | un | ٦. |
|------|-----|----|------|-----|
| Daci | ۸u | ΙU | ui i | ıu. |

Synonyms: Sushi repeat-containing protein SRPX2, SRPX2, Sushi-repeat protein upregulated in

leukemia, SRPUL

Background: Acts as a ligand for the urokinase plasminogen activator surface receptor. Plays a role in angiogenesis by inducing endothelial cell migration and the formation of vascular network (cords). Involved in cellular migration and adhesion. Increases the phosphorylation levels of FAK. Interacts with and increases the mitogenic activity of HGF. Promotes synapse formation. May have a role in the perisylvian region, critical for language and cognitive

Application Details

Application Notes:

IF(IHC-P) 1:50-200

development.

IF(IHC-F) 1:50-200

IF(ICC) 1:50-200

Restrictions:

For Research Use only

Handling

| Format: | Liquid |
|--------------------|--|
| Concentration: | 1 μg/μL |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol. |
| Preservative: | ProClin |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. |
| Storage: | -20 °C |
| Storage Comment: | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles. |
| Expiry Date: | 12 months |