antibodies .- online.com





anti-THO Complex 2 antibody (AA 101-200) (Alexa Fluor 680)



Go to Product page

\sim			
	N/P	r\/	i⊢₩

Quantity:	100 μL
Target:	THO Complex 2 (THOC2)
Binding Specificity:	AA 101-200
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This THO Complex 2 antibody is conjugated to Alexa Fluor 680
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human THOC2
Isotype:	IgG
Cross-Reactivity:	Human
Predicted Reactivity:	Mouse,Rat,Dog,Cow,Pig,Horse,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	THO Complex 2 (THOC2)
Alternative Name:	THOC2 (THOC2 Products)

Target Details

Background:	Synonyms: THO2, CXorf3, hTREX120, dJ506G2.1, THO complex subunit 2, THOC2	
	Background: Required for efficient export of polyadenylated RNA and spliced mRNA. Acts as	
	component of the THO subcomplex of the TREX complex which is thought to couple mRNA	
	transcription, processing and nuclear export, and which specifically associates with spliced	
	mRNA and not with unspliced pre-mRNA. TREX is recruited to spliced mRNAs by a	
	transcription-independent mechanism, binds to mRNA upstream of the exon-junction complex	
	(EJC) and is recruited in a splicing- and cap-dependent manner to a region near the 5' end of the	
	mRNA where it functions in mRNA export to the cytoplasm via the TAP/NFX1 pathway. The	
	TREX complex is essential for the export of Kaposi's sarcoma-associated herpesvirus (KSHV)	
	intronless mRNAs and infectious virus production. THOC2 (and probably the THO complex) is	
	involved in releasing mRNA from nuclear speckle domains. Required for NXF1 localization to	
	the nuclear rim.	
Gene ID:	57187	
UniProt:	Q8NI27	
Application Details		
Application Notes:	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and	
	50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be	
	handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	

		1.	
\vdash	land	lın	
	ıaııu	1111	U

Expiry Date:

12 months