

Datasheet for ABIN5011720

anti-CEMIP antibody (AA 101-200) (AbBy Fluor® 680)



Overview

Overview	
Quantity:	100 μL
Target:	CEMIP
Binding Specificity:	AA 101-200
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CEMIP antibody is conjugated to AbBy Fluor® 680
Application:	Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunofluorescence (Cultured Cells) (IF (cc))
Product Details	
Immunogon:	VIII conjugated cynthetic pontide derived from hyman TMFMOI

Immunogen:	KLH conjugated synthetic peptide derived from human TMEM2L
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Predicted Reactivity:	Dog,Cow,Horse,Chicken,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	CEMIP
Alternative Name:	CEMIP (CEMIP Products)

Target Details

Background:	Synonyms: CCSP1, TMEM2L, KIAA1199, Cell migration-inducing and hyaluronan-binding
	protein, CEMIP, HYBID
	Background: Mediates depolymerization of hyaluronic acid (HA) via the cell membrane-
	associated clathrin-coated pit endocytic pathway. Binds to hyaluronic acid. Hydrolyzes high
	molecular weight hyaluronic acid to produce an intermediate-sized product, a process that may
	occur through rapid vesicle endocytosis and recycling without intracytoplasmic accumulation
	or digestion in lysosomes. Involved in hyaluronan catabolism in the dermis of the skin and
	arthritic synovium. Positively regulates epithelial-mesenchymal transition (EMT), and hence tumor cell growth, invasion and cancer dissemination. In collaboration with HSPA5/BIP,
	promotes cancer cell migration in a calcium and PKC-dependent manner. May be involved in
	hearing.
Gene ID:	57214
UniProt:	Q8WUJ3
Application Details	
Application Notes:	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
	50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
	handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months