antibodies -online.com





anti-CEMIP antibody (AA 101-200) (AbBy Fluor® 750)



Go to Product page

()	۱ ۱	\cap	r	/1	\cap	۱ ۸	1
0	'V	ㄷ	I١	νı	ㄷ	٧	۷

Quantity:	100 μL	
Target:	CEMIP	
Binding Specificity:	AA 101-200	
Reactivity:	Human, Mouse, Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This CEMIP antibody is conjugated to AbBy Fluor® 750	
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human TMEM2L	
Isotype:	IgG	
Cross-Reactivity:	Human, Mouse, Rat	
Predicted Reactivity:	Dog,Cow,Horse,Chicken,Rabbit	
Purification:	Purified by Protein A.	

Target Details

Target:	CEMIP
Alternative Name:	CEMIP (CEMIP Products)

Target Details

Background:	Synonyms: CCSP1, TMEM2L, KIAA1199, Cell migration-inducing and hyaluronan-binding	
	protein, CEMIP, HYBID	
	Background: Mediates depolymerization of hyaluronic acid (HA) via the cell membrane-	
	associated clathrin-coated pit endocytic pathway. Binds to hyaluronic acid. Hydrolyzes high	
	molecular weight hyaluronic acid to produce an intermediate-sized product, a process that may	
	occur through rapid vesicle endocytosis and recycling without intracytoplasmic accumulation	
	or digestion in lysosomes. Involved in hyaluronan catabolism in the dermis of the skin and	
	arthritic synovium. Positively regulates epithelial-mesenchymal transition (EMT), and hence	
	tumor cell growth, invasion and cancer dissemination. In collaboration with HSPA5/BIP,	
	promotes cancer cell migration in a calcium and PKC-dependent manner. May be involved in	
	hearing.	
Gene ID:	57214	
UniProt:	Q8WUJ3	
Application Details		
Application Notes:	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and	
	50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be	
	handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	
Expiry Date:	12 months	