antibodies.com

Datasheet for ABIN5012025 anti-TSEN2 antibody (Alexa Fluor 750)



~ ·	
Overview	

Overview	
Quantity:	100 µL
Target:	TSEN2
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TSEN2 antibody is conjugated to Alexa Fluor 750
Application:	Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p))
Product Details	
Immunogen:	KLH conjugated synthetic peptide derived from human TSEN2
lsotype:	lgG
Cross-Reactivity:	Human, Mouse, Rat
Purification:	Purified by Protein A.
Target Details	
Target:	TSEN2
Alternative Name:	TSEN2 (TSEN2 Products)
Background:	Synonyms: HsSen2, MGC2776, MGC4440, SEN2, SEN2L, tRNA intron endonuclease Sen2, tRNA
	splicing endonuclease 2 homolog SEN2, S. cerevisiae, tRNA splicing endonuclease subunit
	Sen2, TSEN 2, SEN2_HUMAN.
	Background: The tRNA-splicing endonuclease complex is responsible for identifying and

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN5012025 | 03/07/2024 | Copyright antibodies-online. All rights reserved.

Pathways:	Nuclear Receptor Transcription Pathway, Regulation of Lipid Metabolism by PPARalpha
Gene ID:	80746
	epilepsy, extrapyramidal dyskinesia and chorea without spinal cord findings.
	pontocerebellar hypoplasia type 2B, which is characterized by progressive microencephaly with
	active site for 5?-splice site cleavage. Defects in the gene encoding TSEN2 are the cause of
	TSEN2 that are produced as a result of alternative splicing events. Isoform 1 seems to carry the
	catalytic subunits of the tRNA-splicing endonuclease complex. There are three isoforms of
	endonuclease Sen2, is a 465 amino acid nuclear protein that constitutes one of the two
	splicing events. TSEN2 (tRNA-splicing endonuclease subunit Sen2), also known as tRNA-intron
	its association with pre-mRNA processing factors, is thought to link pre-tRNA and pre-mRNA
	splicing, the heterotetrameric endonuclease complex participates in mRNA processing and, via
	molecules with 2',3' cyclic phosphates and 5'-OH termini. In addition to its role in pre-tRNA
	cleaving pre-tRNA at both 5? and 3? splice sites, thereby releasing introns and free tRNA

Application Details

Application Notes:	IF(IHC-P) 1:50-200
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/2 | Product datasheet for ABIN5012025 | 03/07/2024 | Copyright antibodies-online. All rights reserved.