antibodies .- online.com





anti-Taxilin beta antibody (AA 331-430) (Alexa Fluor 750)



Go to Product page

| () | 11/0 | K\ / | iew | 1 |
|-----|--------|--------|---------------------|---|
| | \cup | 'I V/I | $\square \vee \vee$ | ı |
| | | | | |

| Quantity: | 100 μL | |
|----------------------|--|--|
| Target: | Taxilin beta (TXLNB) | |
| Binding Specificity: | AA 331-430 | |
| Reactivity: | Human, Mouse, Rat | |
| Host: | Rabbit | |
| Clonality: | Polyclonal | |
| Conjugate: | This Taxilin beta antibody is conjugated to Alexa Fluor 750 | |
| Application: | Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Western Blotting (WB) | |

Product Details

| Immunogen: | KLH conjugated synthetic peptide derived from human TXLNB |
|-----------------------|---|
| Isotype: | IgG |
| Specificity: | This may have some cross-reactivity with TXLNA and TXLNG. |
| Cross-Reactivity: | Human, Mouse, Rat |
| Predicted Reactivity: | Dog,Cow,Sheep,Pig,Horse |
| Purification: | Purified by Protein A. |

Target Details

| Target: | Taxilin beta (TXLNB) | |
|---------|----------------------|--|

Target Details

| rarger betaile | | | |
|---------------------|---|--|--|
| Alternative Name: | TXLNB (TXLNB Products) | | |
| Background: | Synonyms: MDP77, LST001, C6orf198, dJ522B19.2, Beta-taxilin, Muscle-derived protein 77, | | |
| | hMDP77, TXLNB | | |
| | Background: Beta-taxilin promotes nerve regeneration and may be involved in intracellular | | |
| | vesicle transport. Expressed predominantly in cardiac and skeletal muscle, beta-taxilin binds to | | |
| | the coiled coil region of the syntaxin family members STX1A, STX3A, and STX4A. Beta- and | | |
| | gamma-taxilins, bind to the alpha subunit of the nascent polypeptide-associated complex | | |
| | (NAC) and affect its nuclear distribution, suggesting that the taxilin family is involved not only in | | |
| | the translational process through its interaction with NAC but also in the transcriptional process | | |
| | through its interaction with alphaNAC alone. | | |
| Gene ID: | 167838 | | |
| UniProt: | Q8N3L3 | | |
| Application Details | | | |
| Application Notes: | IF(IHC-P) 1:50-200 | | |
| | IF(IHC-F) 1:50-200 | | |
| | IF(ICC) 1:50-200 | | |
| Restrictions: | For Research Use only | | |
| Handling | | | |
| Format: | Liquid | | |
| Concentration: | 1 μg/μL | | |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and | | |
| | 50 % Glycerol. | | |
| Preservative: | ProClin | | |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be | | |
| | handled by trained staff only. | | |
| Storage: | -20 °C | | |
| Storage Comment: | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles. | | |
| Expiry Date: | 12 months | | |
| | | | |