antibodies -online.com





anti-UFD1L antibody (Alexa Fluor 750)



Go to Product page

| \sim | | | | | |
|--------|----|----|----|-------------|---|
| () | VE | ۲۱ | /1 | \triangle | Λ |

| Quantity: | 100 μL | | |
|-------------------|--|--|--|
| Target: | UFD1L | | |
| Reactivity: | Human, Mouse, Rat | | |
| Host: | Rabbit | | |
| Clonality: | Polyclonal | | |
| Conjugate: | This UFD1L antibody is conjugated to Alexa Fluor 750 | | |
| Application: | Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) | | |
| Product Details | | | |
| Immunogen: | KLH conjugated synthetic peptide derived from human UFD1L | | |
| Isotype: | IgG | | |
| Cross-Reactivity: | Human, Mouse, Rat | | |
| Purification: | Purified by Protein A. | | |
| Target Details | | | |
| Target: | UFD1L | | |
| Alternative Name: | UFD1L (UFD1L Products) | | |
| Background: | Synonyms: UB fusion protein 1, Ubiquitin fusion degradation 1 like yeast, Ubiquitin fusion | | |
| | degradation 1 like, Ubiquitin fusion degradation protein 1 homolog, UFD1, UFD1L. | | |
| | Background: UFD1L is a member of the UFD1 family of proteins and is a component of the | | |

ubiquitin-dependent proteolytic pathway which degrades ubiquitin fusion proteins. This

Target Details

complex, also containing UFD1L, VCP and NPLOC4, binds ubiquitinated proteins and is required for the export of misfolded proteins from the ER to the cytoplasm for disposal. The NPLOC4-UFD1L-VCP complex also regulates spindle disassembly at the end of mitosis and is necessary for the formation of a closed nuclear envelope. UFD1L gene hemizygosity is the cause of some developmental defects including DiGeorge syndrome (DGS), velo-cardio-facial syndrome (VCFS) and Opitz G/BBB syndrome. UFD1L has 2 named isoforms produced by alternative splicing.

Gene ID:

7353

Application Details

| Application Notes: | IF(IHC-P) 1:50-200 |
|--------------------|-----------------------|
| Restrictions: | For Research Use only |
| Handling | |
| Format: | Liquid |
| Concentration: | 1 μg/μL |

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

ProClin Preservative: Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. -20 °C Storage:

Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles. Storage Comment:

Expiry Date: 12 months