antibodies - online.com







anti-UNC5A antibody (AA 125-220) (Alexa Fluor 750)



| () | 1/0 | r\ /1 | 014 | |
|-----|-----|-------|-----|---|
| () | ve | I V I | -v | V |
| | | | | |

| Quantity: | 100 μL | |
|----------------------|---|--|
| Target: | UNC5A | |
| Binding Specificity: | AA 125-220 | |
| Reactivity: | Human | |
| Host: | Rabbit | |
| Clonality: | Polyclonal | |
| Conjugate: | This UNC5A antibody is conjugated to Alexa Fluor 750 | |
| Application: | Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) | |

Product Details

| Immunogen: | KLH conjugated synthetic peptide derived from human Unc5a | |
|-----------------------|---|--|
| Isotype: | IgG | |
| Predicted Reactivity: | Human,Mouse,Rat,Dog,Cow,Sheep,Pig,Horse | |
| Purification: | Purified by Protein A. | |

Target Details

| Target: | UNC5A | |
|--|----------------------------------|--|
| Alternative Name: | e: Unc5a/UNC5H1 (UNC5A Products) | |
| Background: Synonyms: Netrin receptor UNC 5A, Netrin receptor UNC5A, Netrin receptor Unc5h1, Pro | | |

unc-5 homolog 1, Protein unc-5 homolog A, Unc 5 homolog 1, Unc 5 homolog A, Unc 5a, Unc5a, Unc5a, Unc5 homolog 1, Unc5 homolog 1, Unc5 homolog A, Unc5a, UNC5A_HUMAN, Unc5h1, FLJ16449, KIAA1976. Background: The UNC5H family of proteins act as transmembrane receptors for netrin-1 and play a crucial role in axon guidance and migration of neural cells. Additionally, when cleaved by a caspase to produce an intracellular fragment containing a death domain,UNC5H receptors induce apoptosis. This activity is blocked by the binding of netrin-1. In the absence of netrin-1, UNC5H receptors act as tumor suppressors by inhibiting anchorage-independent growth and invasion, but mutation of these receptors provides a potential mechanism for tumorigenicity. The expression of UNC5H receptors is downregulated in multiple carcinomas, including colorectal, breast, ovary, uterus, stomach, lung and kidney cancers. UNC5H1, also known as UNC5HA (unc-5 homolog A), is a member of the UNC5H family of proteins and is localized to the cell membrane. Three isoforms of UNC5H1 exist due to alternative splicing events.

Application Details

| Application Notes: | IF(IHC-P) 1:50-200 |
|--------------------|-----------------------|
| | IF(IHC-F) 1:50-200 |
| | IF(ICC) 1:50-200 |
| Restrictions: | For Research Use only |

Handling

| Format: | Liquid | |
|--------------------|--|--|
| Concentration: | 1 μg/μL | |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol. | |
| Preservative: | ProClin | |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. | |
| Storage: | -20 °C | |
| Storage Comment: | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles. | |
| Expiry Date: | 12 months | |