

[Go to Product page](#)

Datasheet for ABIN5012677

**anti-ZFYVE27 antibody (AA 341-411) (Alexa Fluor 750)**

## Overview

Quantity:	100 µL
Target:	ZFYVE27
Binding Specificity:	AA 341-411
Reactivity:	Human, Rat, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ZFYVE27 antibody is conjugated to Alexa Fluor 750
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

## Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human ZFYVE27
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Predicted Reactivity:	Dog,Cow,Rabbit
Purification:	Purified by Protein A.

## Target Details

Target:	ZFYVE27
Alternative Name:	ZFYVE27 ( <a href="#">ZFYVE27 Products</a> )

## Target Details

Background:	<p>Synonyms: Protrudin, RP11 459F3.2, SPG33, ZFY27_HUMAN, ZFYVE27, zinc finger FYVE domain containing 27, Zinc finger FYVE domain containing protein 27, Zinc finger FYVE domain-containing protein 27.</p> <p>Background: Zinc finger FYVE domain-containing protein 27 (ZFYVE27), also known as SPG33, is a 411 amino acid member of the FYVE-finger family of proteins. The FYVE domain is a cysteine-rich domain of about 70 amino acids that plays a role in the endosomal localization of the FYVE-finger proteins, and a majority of these proteins serve as regulators of endocytic membrane trafficking. ZFYVE27, a multi-pass membrane protein, is an endosomal protein that binds to Spastin, a protein that is primarily involved in microtubule dynamics and severing, vesicular trafficking and endosomal trafficking. Mutations in the gene encoding ZFTVE27 affect neuronal intracellular trafficking in the corticospinal tract and are thought to lead to hereditary spastic paraplegia (HSP), a neurodegenerative disorder, characterized by progressive paralysis of the legs, which is caused by impaired axonal transport. Five isoforms of ZFYVE27 exist as a result of alternative splicing events.</p>
Gene ID:	118813
UniProt:	<a href="#">Q5T4F4</a>
Pathways:	<a href="#">Neurotrophin Signaling Pathway</a>

## Application Details

Application Notes:	IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200
Restrictions:	For Research Use only

## Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS ( pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

## Handling

---

Storage:	-20 °C
----------	--------

Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
------------------	---

Expiry Date:	12 months
--------------	-----------