

Datasheet for ABIN5013246 anti-BCR antibody (AA 1042-1268)

2 Images

Background:




Go to Product page

Overview	
Quantity:	100 µL
Target:	BCR
Binding Specificity:	AA 1042-1268
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This BCR antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC)
Product Details	
lmmunogen:	BCR (Met1042-Thr1268)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against BCR. It has been selected for its
	ability to recognize BCR in immunohistochemical staining and western blotting.
Purification:	Antigen-specific affinity chromatography
Target Details	
Target:	BCR
Abstract:	BCR Products
D l l -	All II ALL DOD/FOFD4 DOD4 OAN FOFD4/DOD DUI D. I

Alternative Names: ALL, BCR/FGFR1, BCR1, CML, FGFR1/BCR, PHL, Renal carcinoma antigen


Application Details • Western blotting: 1:50-400 Immunocytochemistry in formalin fixed cells: 1:50-500 Immunohistochemistry in formalin fixed frozen section: 1:50-500 Immunohistochemistry in paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-1:5000 Optimal working dilutions must be determined by end user. Comment: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious		
Application Details - Western blotting: 1:50-400 Immunocytochemistry in formalin fixed cells: 1:50-500 Immunohistochemistry in formalin fixed frozen section: 1:50-500 Immunohistochemistry in paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-1:5000 Optimal working dilutions must be determined by end user. Comment: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 378degC for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition. Restrictions: For Research Use only Handling Format: Liquid Concentration: Lot specific		NY-REN-26
Application Notes: • Western blotting: 1:50-400 Immunocytochemistry in formalin fixed cells: 1:50-500 Immunohistochemistry in formalin fixed frozen section: 1:50-500 Immunohistochemistry in paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-1:5000 Optimal working dilutions must be determined by end user. Comment: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition. Restrictions: For Research Use only Handling Format: Liquid Concentration: Lot specific	Pathways:	Regulation of Leukocyte Mediated Immunity, Platelet-derived growth Factor Receptor Signaling
Immunohistochemistry in formalin fixed frozen section: 1:50-500 Immunohistochemistry in paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-1:5000 Optimal working dilutions must be determined by end user. Comment: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition. Restrictions: For Research Use only Handling Format: Liquid Concentration: Lot specific	Application Details	
thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition. Restrictions: For Research Use only Handling Format: Liquid Concentration: Lot specific	Application Notes:	Immunohistochemistry in formalin fixed frozen section: 1:50-500 Immunohistochemistry in paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-1:5000 Optimal
Handling Format: Liquid Concentration: Lot specific	Comment:	degradation and precipitation were observed. The loss rate is less than 5% within the expiration
Format: Liquid Concentration: Lot specific	Restrictions:	For Research Use only
Concentration: Lot specific	Handling	
	Format:	Liquid
Buffer: PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.	Concentration:	Lot specific
	Buffer:	PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.

Images



Western Blotting

Image 1. Figure. Western Blot; Sample: Recombinant protein.



Immunohistochemistry

Image 2. Used in DAB staining on fromalin fixed paraffinembedded liver tissue