antibodies -online.com





anti-Calneuron 1 antibody (AA 1-192)





| Overview | |
|----------------------|--|
| Quantity: | 100 μL |
| Target: | Calneuron 1 (CALN1) |
| Binding Specificity: | AA 1-192 |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This Calneuron 1 antibody is un-conjugated |
| Application: | Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC) |
| Product Details | |
| Immunogen: | CALN1 (Met1-Ser192) |

| Immunogen: | CALN1 (Met1-Ser192) |
|---------------|--|
| Isotype: | IgG |
| Specificity: | The antibody is a rabbit polyclonal antibody raised against CALN1. It has been selected for its ability to recognize CALN1 in immunohistochemical staining and western blotting. |
| Purification: | Antigen-specific affinity chromatography |


Target Details

| Target: | Calneuron 1 (CALN1) |
|-------------|---|
| Abstract: | CALN1 Products |
| Background: | Alternative Names: CABP8, Calcium-Binding Protein 8 |

Application Details


| Application Notes: | Western blotting: 1:50-400 Immunocytochemistry in formalin fixed cells: 1:50-500 Immunohistochemistry in formalin fixed frozen section: 1:50-500 Immunohistochemistry in paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-1:5000 Optimal working dilutions must be determined by end user. |
|--------------------|--|
| Comment: | The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition. |
| Restrictions: | For Research Use only |
| Handling | |
| Format: | Liquid |
| Concentration: | Lot specific |
| Buffer: | PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol. |

Images



Western Blotting

Image 1. Figure. Western Blot; Sample: Recombinant protein.



Immunohistochemistry

Image 2. Used in DAB staining on fromalin fixed paraffinembedded Kidney tissue