antibodies -online.com





anti-CPN2 antibody (AA 249-540)

2 Images



| _ | | |
|-------|------|--|
| ()Ver | view | |

| Quantity: | 100 μL |
|--|---|
| Target: | CPN2 |
| Binding Specificity: | AA 249-540 |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This CPN2 antibody is un-conjugated |
| Application: | Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC) |
| Product Details | |
| | |
| Immunogen: | CPN2 (GIn249-Ala540) |
| Immunogen: Isotype: | CPN2 (Gln249-Ala540) IgG |
| | |
| Isotype: | IgG The antibody is a rabbit polyclonal antibody raised against CPN2. It has been selected for its |
| Isotype: Specificity: | IgG The antibody is a rabbit polyclonal antibody raised against CPN2. It has been selected for its ability to recognize CPN2 in immunohistochemical staining and western blotting. |
| Isotype: Specificity: Purification: | IgG The antibody is a rabbit polyclonal antibody raised against CPN2. It has been selected for its ability to recognize CPN2 in immunohistochemical staining and western blotting. |
| Isotype: Specificity: Purification: Target Details | IgG The antibody is a rabbit polyclonal antibody raised against CPN2. It has been selected for its ability to recognize CPN2 in immunohistochemical staining and western blotting. Antigen-specific affinity chromatography |

Carboxypeptidase N regulatory subunit, Carboxypeptidase N polypeptide 2

Application Details

Application Notes:

Western blotting: 1:50-400 Immunocytochemistry in formalin fixed cells: 1:50-500
 Immunohistochemistry in formalin fixed frozen section: 1:50-500 Immunohistochemistry in paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-1:5000 Optimal working dilutions must be determined by end user.

Comment:

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.


Restrictions:

For Research Use only

Handling


| Format: | Liquid |
|----------------|---|
| Concentration: | Lot specific |
| Buffer: | PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol. |

Images



Western Blotting

Image 1. Figure. Western Blot; Sample: Recombinant protein.



Immunohistochemistry

Image 2. Used in DAB staining on fromalin fixed paraffinembedded stomach tissue