

Datasheet for ABIN5013296  
**anti-SLC25A20 antibody (AA 1-301)**[Go to Product page](#)

## 2 Images

## Overview

Quantity:	100 µL
Target:	SLC25A20
Binding Specificity:	AA 1-301
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This SLC25A20 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC)

## Product Details

Immunogen:	CACT (Met1-Leu301)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against CACT. It has been selected for its ability to recognize CACT in immunohistochemical staining and western blotting.
Purification:	Antigen-specific affinity chromatography

## Target Details

Target:	SLC25A20
Alternative Name:	Carnitine Acylcarnitine Translocase (CACT) ( <a href="#">SLC25A20 Products</a> )
Background:	Alternative Names: SLC25A20, CAC, Solute Carrier Family 25 Member 20, Carnitine

## Target Details

Acylcarnitine Carrier, Mitochondrial carnitine/acylcarnitine carrier protein

Pathways: [TCR Signaling](#), [Production of Molecular Mediator of Immune Response](#), [Maintenance of Protein Location](#), [Toll-Like Receptors Cascades](#)


## Application Details

Application Notes:	<ul style="list-style-type: none"><li>Western blotting: 1:50-400 Immunocytochemistry in formalin fixed cells: 1:50-500 Immunohistochemistry in formalin fixed frozen section: 1:50-500 Immunohistochemistry in paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-1:5000 Optimal working dilutions must be determined by end user.</li></ul>
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37&degC for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.
Restrictions:	For Research Use only

## Handling


Format:	Liquid
Concentration:	Lot specific
Buffer:	PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.

## Images



### Western Blotting

**Image 1.** Figure. Western Blot; Sample: Recombinant protein.



#### Immunohistochemistry

**Image 2.** Used in DAB staining on formalin fixed paraffin-embedded Kidney tissue