# antibodies -online.com







# anti-CCRK antibody (AA 4-288)

**Images** 



$\sim$					
()	VE	۲۱	/1	$\triangle$	Λ

Overview	
Quantity:	100 μL
Target:	CCRK (CDK20)
Binding Specificity:	AA 4-288
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CCRK antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC)
Product Details	
Immunogon:	CDK20 (Tyr/LDbo288)

Immunogen:	CDK20 (Tyr4-Phe288)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against CDK20. It has been selected for its ability to recognize CDK20 in immunohistochemical staining and western blotting.
Purification:	Antigen-specific affinity chromatography

# **Target Details**

Target:	CCRK (CDK20)
Alternative Name:	Cyclin Dependent Kinase 20 (CDK20) (CDK20 Products)
Background: Alternative Names: CCRK, CDCH, p42, Cell Cycle Related Kinase, CDK-activating kinase	

Cyclin-dependent protein kinase H, Cyclin-kinase-activating kinase p42

# **Application Details**

#### Application Notes:

Western blotting: 1:50-400 Immunocytochemistry in formalin fixed cells: 1:50-500
 Immunohistochemistry in formalin fixed frozen section: 1:50-500 Immunohistochemistry in paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-1:5000 Optimal working dilutions must be determined by end user.

#### Comment:

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37&degC for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.


Restrictions:

For Research Use only

# Handling


Format:	Liquid
Concentration:	Lot specific
Buffer:	PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.

### **Images**



## **Western Blotting**

**Image 1.** Figure. Western Blot; Sample: Recombinant protein.



# **Immunohistochemistry**

**Image 2.** Used in DAB staining on fromalin fixed paraffinembedded heart tissue