antibodies -online.com





anti-Cytohesin 4 antibody (AA 1-379)

3 Images



Overview

Overview	
Quantity:	100 μL
Target:	Cytohesin 4 (CYTH4)
Binding Specificity:	AA 1-379
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Cytohesin 4 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC)
Product Details	

Immunogen:	CYTH4 (Met1-Phe379)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against CYTH4. It has been selected for its ability to recognize CYTH4 in immunohistochemical staining and western blotting.
Purification:	Antigen-specific affinity chromatography

Target Details

Target:	Cytohesin 4 (CYTH4)
Abstract:	CYTH4 Products
Background:	Alternative Names: PSCD4, CYT4, Pleckstrin Homology, Sec7 And Coiled-Coil Domains 4


Application Details

Application Notes:	Western blotting: 1:50-400 Immunocytochemistry in formalin fixed cells: 1:50-500 Immunohistochemistry in formalin fixed frozen section: 1:50-500 Immunohistochemistry in paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-1:5000 Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	Lot specific
Buffer:	PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.
Images	




Western Blotting

Image 1. Western Blot; Sample: Porcine Liver lysate; Primary Ab: 3μg/ml Rabbit Anti-Human CYTH4 Antibody Second Ab: 0.2μg/mL HRP-Linked Caprine Anti-Rabbit IgG Polyclonal Antibody (Catalog: SAA544Rb19)



Western Blotting

Image 2. Figure. Western Blot; Sample: Recombinant protein.



Immunohistochemistry

Image 3. Used in DAB staining on fromalin fixed paraffinembedded Kidney tissue