antibodies - online.com





Datasheet for ABIN5013459

anti-DPP7 antibody (AA 28-486)





| _ | | | | | |
|---|---|----|----|---|---|
| U | V | er | VI | е | W |

Alternative Name:

Background:

| Quantity: | 100 μL | |
|----------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--|
| Target: | DPP7 | |
| Binding Specificity: | AA 28-486 | |
| Reactivity: | Human | |
| Host: | Rabbit | |
| Clonality: | Polyclonal | |
| Conjugate: | This DPP7 antibody is un-conjugated | |
| Application: | Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC) | |
| | | |
| Product Details | | |
| Product Details Immunogen: | DPP7 (Asp28-Gly486) | |
| | DPP7 (Asp28-Gly486) IgG | |
| Immunogen: | IgG The antibody is a rabbit polyclonal antibody raised against DPP7. It has been selected for its | |
| Immunogen: Isotype: | IgG | |
| Immunogen: Isotype: Specificity: | IgG The antibody is a rabbit polyclonal antibody raised against DPP7. It has been selected for its ability to recognize DPP7 in immunohistochemical staining and western blotting. | |


Alternative Names: DPP2, DPPII, QPP, Dipeptidylpeptidase 7

Dipeptidyl Peptidase 7 (DPP7) (DPP7 Products)

Application Details


| Application Notes: | Western blotting: 1:50-400 Immunocytochemistry in formalin fixed cells: 1:50-500 Immunohistochemistry in formalin fixed frozen section: 1:50-500 Immunohistochemistry in paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-1:5000 Optimal working dilutions must be determined by end user. | |
|--------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--|
| Comment: | The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition. | |
| Restrictions: | For Research Use only | |
| Handling | | |
| Format: | Liquid | |
| Concentration: | Lot specific | |
| Buffer: | PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol. | |

Images




Western Blotting

Image 1. Figure. Western Blot; Sample: Recombinant protein.



Immunohistochemistry

Image 2. DAB staining on IHC-P; Samples: Human Stomach
Tissue



Immunohistochemistry

Image 3. Used in DAB staining on fromalin fixed paraffinembedded Kidney tissue

Please check the product details page for more images. Overall 4 images are available for ABIN5013459.