antibodies .- online.com





anti-DOK3 antibody (AA 45-310)

3 Images



Go to Product page

\sim					
	1//6	r	V I	Θ	Λ

- OVERVIEW		
Quantity:	100 μL	
Target:	DOK3	
Binding Specificity:	AA 45-310	
Reactivity:	Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This DOK3 antibody is un-conjugated	
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC)	
Product Details		
i Toduct Details		
Immunogen:	DOK3 (Ser45-Pro310)	
	DOK3 (Ser45-Pro310) IgG	
Immunogen:		
Immunogen: Isotype:	IgG The antibody is a rabbit polyclonal antibody raised against DOK3. It has been selected for its	
Immunogen: Isotype: Specificity:	IgG The antibody is a rabbit polyclonal antibody raised against DOK3. It has been selected for its ability to recognize DOK3 in immunohistochemical staining and western blotting.	
Immunogen: Isotype: Specificity: Purification:	IgG The antibody is a rabbit polyclonal antibody raised against DOK3. It has been selected for its ability to recognize DOK3 in immunohistochemical staining and western blotting.	
Immunogen: Isotype: Specificity: Purification: Target Details	IgG The antibody is a rabbit polyclonal antibody raised against DOK3. It has been selected for its ability to recognize DOK3 in immunohistochemical staining and western blotting. Antigen-specific affinity chromatography	


Application Details

Application Notes:	 Western blotting: 1:50-400 Immunocytochemistry in formalin fixed cells: 1:50-500 Immunohistochemistry in formalin fixed frozen section: 1:50-500 Immunohistochemistry in paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-1:5000 Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	Lot specific
Buffer:	PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.
lmages	




Western Blotting

Image 1. Western Blot; Sample: Rat Liver lysate; Primary Ab: 2μg/ml Rabbit Anti-Rat DOK3 Antibody Second Ab: 0.2μg/mL HRP-Linked Caprine Anti-Rabbit IgG Polyclonal Antibody (Catalog: SAA544Rb19)



Western Blotting

Image 2. Figure. Western Blot; Sample: Recombinant protein.



Immunohistochemistry

Image 3. Used in DAB staining on fromalin fixed paraffinembedded Kidney tissue