
antibodies -online.com







Images




Go to Product page

_					
U	V	er	VI	е	W

Quantity:	100 μL	
Target:	DTNBP1	
Binding Specificity:	AA 110-305	
Reactivity:	Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This DTNBP1 antibody is un-conjugated	
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC)	
Product Details		
Immunogen:	DTNBP1 (Asp110-Leu305)	
Isotype:	IgG	
Specificity:	The antibody is a rabbit polyclonal antibody raised against DTNBP1. It has been selected for its ability to recognize DTNBP1 in immunohistochemical staining and western blotting.	
Purification:	Antigen-specific affinity chromatography	
Target Details		
Target:	DTNBP1	
Abstract:	DTNBP1 Products	
Background:	Alternative Names: DBND, HPS7, My031, SDY, Dysbindin, Biogenesis of lysosome-related	


paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-1:5000 Optimal working dilutions must be determined by end user. Comment: The thermal stability is described by the loss rate. The loss rate was determined by accele thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious			
Application Details - Western blotting: 1:50-400 Immunocytochemistry in formalin fixed cells: 1:50-500 Immunohistochemistry in formalin fixed frozen section: 1:50-500 Immunohistochemistry paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-1:5000 Optimal working dilutions must be determined by end user. Comment: The thermal stability is described by the loss rate. The loss rate was determined by accele thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expir date under appropriate storage condition. Restrictions: For Research Use only Handling Format: Liquid Concentration: Lot specific		organelles complex 1 subunit 8, Hermansky-Pudlak syndrome 7 protein	
Application Notes: • Western blotting: 1:50-400 Immunocytochemistry in formalin fixed cells: 1:50-500 Immunohistochemistry in formalin fixed frozen section: 1:50-500 Immunohistochemistry paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-1:5000 Optimal working dilutions must be determined by end user. Comment: The thermal stability is described by the loss rate. The loss rate was determined by accele thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiritate under appropriate storage condition. Restrictions: For Research Use only Handling Format: Liquid Concentration: Lot specific	Pathways:	Synaptic Membrane, Regulation of G-Protein Coupled Receptor Protein Signaling	
Immunohistochemistry in formalin fixed frozen section: 1:50-500 Immunohistochemistry paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-1:5000 Optimal working dilutions must be determined by end user. Comment: The thermal stability is described by the loss rate. The loss rate was determined by acceled thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiral date under appropriate storage condition. Restrictions: For Research Use only Handling Format: Liquid Concentration: Lot specific	Application Details		
thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expir date under appropriate storage condition. Restrictions: For Research Use only Handling Format: Liquid Concentration: Lot specific	Application Notes:	Immunohistochemistry in formalin fixed frozen section: 1:50-500 Immunohistochemistry in paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-1:5000 Optimal	
Handling Format: Liquid Concentration: Lot specific	Comment:	degradation and precipitation were observed. The loss rate is less than 5% within the expiration	
Format: Liquid Concentration: Lot specific	Restrictions:	For Research Use only	
Concentration: Lot specific	Handling		
<u>'</u>	Format:	Liquid	
Buffer: PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.	Concentration:	Lot specific	
	Buffer:	PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.	






Western Blotting

Image 1. Western Blot; Sample: Rat Cerebrum lysate; Primary Ab: 2μg/ml Rabbit Anti-Rat DTNBP1 Antibody Second Ab: 0.2μg/mL HRP-Linked Caprine Anti-Rabbit IgG Polyclonal Antibody (Catalog: SAA544Rb19)



Western Blotting

Image 2. Figure. Western Blot; Sample: Recombinant protein.



Immunohistochemistry

Image 3. Used in DAB staining on fromalin fixed paraffinembedded spleen tissue