antibodies -online.com







anti-EIF2D antibody (AA 1-117)

2 Images




\sim					
	1//	⊃r	V/I	Φ\	Λ

OVCIVICVV		
Quantity:	100 μL	
Target:	EIF2D	
Binding Specificity:	AA 1-117	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This EIF2D antibody is un-conjugated	
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC)	
Product Details		
Immunogen:	EIF2D (Met1-Leu117)	
Isotype:	IgG	
Specificity:	The antibody is a rabbit polyclonal antibody raised against EIF2D. It has been selected for its ability to recognize EIF2D in immunohistochemical staining and western blotting.	
Purification:	Antigen-specific affinity chromatography	
Target Details		
Target:	EIF2D	
Abstract:	EIF2D Products	
Background:	Alternative Names: HCA56, LGTN, Hepatocellular carcinoma-associated antigen 56, Ligatin	

Application Details


Application Notes:	Western blotting: 1:50-400 Immunocytochemistry in formalin fixed cells: 1:50-500 Immunohistochemistry in formalin fixed frozen section: 1:50-500 Immunohistochemistry in paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-1:5000 Optimal working dilutions must be determined by end user.	
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	Lot specific	
Buffer:	PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.	

Images



Western Blotting

Image 1. Figure. Western Blot; Sample: Recombinant protein.



Immunohistochemistry

Image 2. Used in DAB staining on fromalin fixed paraffinembedded Kidney tissue