antibodies - online.com







anti-ERI1 antibody (AA 2-249)





\sim					
	1//	r۱.	/ I	\triangle	٨

Overview	
Quantity:	100 μL
Target:	ERI1
Binding Specificity:	AA 2-249
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ERI1 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC)
Product Details	
Immunogen:	ERI1 (Glu2-Lys249)

Immunogen:	ERI1 (Glu2-Lys249)	
Isotype:	IgG	
Specificity:	The antibody is a rabbit polyclonal antibody raised against ERI1. It has been selected for its ability to recognize ERI1 in immunohistochemical staining and western blotting.	
Purification:	cation: Antigen-specific affinity chromatography	

Target Details

Target:	ERI1
Abstract:	ERI1 Products
Background:	Alternative Names: THEX1, 3'HEX0, 3'-5' exonuclease ERI1, Histone mRNA 3'-end-specific

exoribonuclease, Enhanced RNAi Three Prime mRNA Exonuclease Homolog 1

Application Details

Application Notes:

Western blotting: 1:50-400 Immunocytochemistry in formalin fixed cells: 1:50-500
 Immunohistochemistry in formalin fixed frozen section: 1:50-500 Immunohistochemistry in paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-1:5000 Optimal working dilutions must be determined by end user.

Comment:

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.


Restrictions:

For Research Use only

Handling


Format:	Liquid
Concentration:	Lot specific
Buffer: PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.	

Images



Western Blotting

Image 1. Figure. Western Blot; Sample: Recombinant protein.



Immunohistochemistry

Image 2. Used in DAB staining on fromalin fixed paraffinembedded kidney tissue